

**JIMMA UNIVERSITY
COLLEGE OF NATURAL SCIENCES
SCHOOL OF GRADUATE STUDIES
DEPARTMENT OF CHEMISTRY**



M. Sc THESIS

ON

**PHYTOCHEMICAL INVESTIGATION OF *Dodonaea angustifolia* LEAVES AND
EVALUATIO OF ITS ANTIBACTERIAL ACTIVITIES**

OCTOBER, 2017
JIMMA, ETHIOPIA

PHYTOCHEMICAL INVESTIGATION OF *DONAEA ANGUSTIFOLIA* LEAVES AND
EVALUATION OF ITS ANTIBACTERIAL ACTIVITIES

A THESIS SUBMITTED TO SCHOOL OF GRADUATE STUDIES JIMMA
UNIVERSITY IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE
DEGREE OF MASTERS OF SCIENCE IN CHEMISTRY

BY

GETACHEW NEMERA URGESA

ADVISORS

TSEGAYE DEYOU (PhD)

AND

NEGERA ABDISSA (PhD)

OCTOBER, 2017

Jimma University
School of Graduate studies

Phytochemical Investigation of *Dodonaea angustifolia* Leaves and Evaluation of its
Antibacterial Activities

By
Getachew Namera Urgessa

A Thesis Submitted to School of Graduate Studies Jimma University in Partial Fulfillment of
the Requirements for the Degree of Master of Science in Organic Chemistry

Approved by Board of Examiners

External Examiner	Signature	Date
_____	_____	_____
Internal Examiner		
_____	_____	_____
Advisors		
_____	_____	_____
_____	_____	_____
Head, Department of Chemistry		
_____	_____	_____

Declaration

I declare that this thesis is my original work and has not been presented anywhere for award of any degree or diploma in any University.

Getachew Namera

This M.Sc. Thesis has been submitted with our approval as supervisors:

Tsegaye Deyou (PhD)
Chemistry Department
Jimma University

Negera Abdissa (PhD)
Chemistry Department
Jimma University

Table of Contents

Table of Contents	page
Table of Contents	i
List of Figures	iv
Lists of Appendices.....	v
List of Abbreviations	vi
Acknowledgements.....	vii
Abstract.....	viii
1. Introduction	1
1.1. Background of the Study.....	1
1.2. Objectives of the Study	3
1.3.1. General Objective	3
1.3.2. Specific Objectives	3
2. Review of Related Literature	5
2.1. The Microbial Burden	5
2.1.1 Economical Burden of Infection Disease	7
2.2.2 Traditional Uses of the Genus <i>Dodonaea</i>	11
2.2.3 Compounds from Genus <i>Dodonaea</i>	11
2.2.4. Biological Activities of the Genus <i>Dodonaea</i>	14
3 Materials and Methods.....	15
3.1 Chemical, Apparatus and Equipment.....	15
3.2 Collection and Preparation of Plant specimens.....	15
3.3 Extraction and Isolation	16
3.3.1 Extraction of Leaves of <i>Dodonaea angustifolia</i>	16

3.3.2 Isolation of Leaves of <i>Dodonaea angustifolia</i> extract.....	16
3.4. Culture Media.....	18
3.4.1. Test Strains.....	18
3.4.2 Standardization of Inoculums.....	18
3.5 Antibacterial Activity Test.....	19
3.5.1 Agar Disc Diffusion Method.....	19
4. Results and Discussion.....	20
4.1 Secondary Metabolites from leaves of <i>Dodonaea angustifolia</i>	20
4.2 Characterization of Isolated Compound 1a.....	20
2.1. Summary of Spectra and Physical data the Isolated Compound (1a).	25
4.3 Biological Assay.....	26
4.3.1 Antibacterial activities of the extracts and isolated compound of the leaves of <i>Dodonaea angustifolia</i>	26
5. Conclusion and Recommendation.....	28
5.1 Conclusion.....	28
5.2 Recommendation.....	29
6. Reference.....	30
7. Appendices.....	36

Lists of Table

List of table	page
Table 1: List of compounds isolated from <i>Dodonaea</i> species.....	11
Table 2: The ^1H and ^{13}C NMR spectral data of compound (1A) and literature for santin (43).....	21
Table 3. HSQC data for compound 1A.	22
Table 4. Observed correlations in HMBC data of compound 1A	23
Table 5. Antibacterial activity of <i>Dodonaea angustifolia</i> leaves crude $\text{CHCl}_3/\text{CH}_3\text{OH}$, ethyl acetate soluble extracts and isolated compound	26

List of Figures

List of figures	page
Figure. 1. Structure of some isolated Drug from plant.	1
Figure 2. structure some plant based compound for antibacterial	6
Figure 3. A picture of <i>Dodonaea angustifolia</i> on uncultivated area taken from Estern Wollega ,Gobu Sayo woreda, Anno district in 2016.	10
Figure 5 Partial structure of compound 1A based on Cosy spectrum.....	22
Figure 6. Partial structure of compound 1A based on HMBC spectra	23
Figure 7. Partial structure of compound 1A based on HMBC spectra	24
Figure 8. Partial structure compound 1A based on HMBC spectra.....	24
Figure 9. Partial structure of compound 1A based on HMBC spectra	24
Figure 10. Final Structure of Isolated compound 1A.	25

Lists of Appendices

Appendix	page
Appendix 1. Bioassay tests of crude extracts zone of growth inhibition.....	39
Appendix 2. Antibacterial activities of different concentrations of ethyl acetate soluble extracts.....	39
Appendix 3. Bioassays tests of isolated compounds' zone of growth inhibition	39
Appendix 4. ^1H NMR spectrum of compound 1A in CDCl_3	40
Appendix 5. ^{13}C NMR spectrum of compound 1A in CDCl_3	41
Appendix 6. COSY spectrum of compound 1A in CDCl_3	42
Appendix 7. HSQC spectrum of compound 1A in CDCl_3	44
Appendix 8. HMBC spectrum of compound 1A in CDCl_3	47

List of Abbreviations

ATCC	American Type Culture Collection
CC	Column Chromatography
COSY	Correlation Spectroscopy
HSQC	Heteronuclear Single Quantum Correlation
HMBC	Heteronuclear Multiple-Bond Correlation
MHB	Mueller-Hinton Broth
MIC	Minimum Inhibitory Concentration
NMR	Nuclear Magnetic Resonance
STDs	Sexually Transmitted Diseases
TLC	Thin-layer Chromatography
TMS	Tetramethylsilane
UV	Ultraviolet
WHO	World Health Organization
HIV	Human Immunodeficiency Virus
AIDS	Acquired Immunodeficiency Syndrome

Acknowledgements

First of all, I would like to thank the Almighty God who gives me full health, peace, wisdom, and acquaintance to realize this thesis work and for still be on my side forever.

I am very much grateful to my advisors; Dr. Tsegaye Deyou and Dr. Negera Abdissa for their remarkable professional support during the course of this study. Their constructive and critical feedbacks added a great value to this thesis work.

It is also my great pleasure to thank staff member of the Department of Chemistry and Department of Biology, Jimma University for their support and cooperation throughout the study period. My special thank goes to my friends; Mr. Dele Abdisa, Mr. Melaku Meshasha, Mr. Melkamu Feyera and Mr. Abebe Diro for their unconditional love, moral, financial backing and sharing my personal burdens. I truly appreciate the encouragement and thoughtful attitudes you all accorded me during my stay at Jimma University. I am also highly grateful to my family, especially my mam, Mrs. Alemitu Yadeta, for her unconditional support, heartfelt moral and love she has accorded me in due course of my academic life since my early schooling to this level. Her unreserved support at different stage of my academic life had a significant effort on the success of my present day study.

Lastly, I would like to express my deepest gratitude to my wife Dule Teferi for her understanding and patience during my absence for the study.

Abstract

Natural products from plants and other natural sources have been providing noble and clinically active drugs. The plant-based traditional medicinal system continues to play significant role in health care, in which approximately 80% of the world's population still rely mainly on traditional medicines for their primary health-care. Therefore, this study was undertaken to identify bioactive secondary metabolites from the leaves of *Dodonaea angustifolia* for antibacterial activities. With this regard, the air dried leaves of *Dodonaea angustifolia* were extracted successively with chloroform/methanol (1:1) by maceration. The crude extract chloroform/methanol extract was then partitioned between Ethyl acetate and H₂O to give Ethyl acetate extract. Both CHCl₃/CH₃OH and ethyl acetate crude extracts were evaluated for their antibacterial activity against four bacterial strains, namely; *Staphylococcus aureus* (ATCC 25923), *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853) and *Enterococcus faecalis* (ATCC 29212). The crude extracts showed antibacterial activities with zone of inhibition ranging from 6-22 mm. The observed activity against *S. aureus* (zone of inhibition, 22 mm) for ethyl acetate soluble extract was comparable with that of the reference drug, gentamycin, which displayed 23 mm zone of inhibition on the same strain. Based on the superior activity of the ethyl acetate extract, it was subjected to fractionation using column chromatography (CC) over silica gel eluted with petroleum ether containing increasing amounts of ethyl acetate, which resulted with the identification of 5,7-dihydroxy-3,4,6-trimethoxyflavone (**1A**) which showed moderate antibacterial activity against the tested strains (*E. faecalis*, *E. coli*, *S. aureus*, and *Ps. aeruginosa*). The structure of the isolated compound was established using 1D (¹H, ¹³C) and 2D NMR (COSY, HSQC and HMBC) spectroscopic techniques. Devising alternative method of extraction as well as carrying the antioxidant, antifungal and antiplasmodium activities of this plant are recommended for further researchs

1. Introduction

1.1. Background of the Study

The importance of medicinal plants in traditional health care system have been started long time ago and still continue to serve the 80% world populations for their primary healthcare [1, 2]. The traditional usage of medicinal plants is more common in developing countries including Africa, Asia and South America. It is also true that the use of traditional medicine is not completely omitted in developed countries as public interest in natural therapies has increased greatly in industrialized countries. For example, in the United States, about 38% of adults and 12% of children were using some form of traditional medicine in the year, 2007 [3]. In addition to the traditional uses, plants have been also serving as a source of modern drugs in the treatment of malaria by quinine (**1**) from *Cinchona* bark and artemisinin (**2**) [4] from *Artemisia annua*, cancer by the vinca alkaloids, vinblastine (**3**) and vincristine (**4**) from the Madagascan periwinkle, *Catharantus roseus* [5] and taxol (**5**) from some Yew species, and for other infectious diseases




Figure. 1. Structure of some isolated Drug from plant.

In general, there is a trend in the world today to shift to natural substances due to various side effects associated to some synthetic drugs with about 200 plant-derived chemical compounds of known structures are being used as drugs or as agents that lead to improvement of human health [6]. During the last two decades, the development of drug resistance as well as the appearance of undesirable side effects of certain antibiotics has led to the search for new antimicrobial agents mainly from plant extracts with the goal to discover new chemical structures [7, 8]. There have been also increased attentions towards finding natural occurring antioxidants to replace synthetic antioxidants which are being restricted due to their carcinogenicity [9].

Like the rest part of the world, the uses of plants as a medicine is also common in Ethiopia, with close to 80% of the population is still dependent on traditional medicine [10]. However, the work that has been done to investigate Ethiopian traditional medicinal plants both phytochemically and biologically is too scanty. Therefore, the current study is focused on phytochemical investigation of one of the Ethiopian Medicinal plants, *Dodonaea angustifolia* for antibacterial principles that could replace the current drugs that are becoming ineffective.

1.2 Statement of the Problem

Infectious diseases are one of the major causes of several numbers of deaths of children, young and adults in developing countries [11]. It is the second leading cause of death worldwide, and the third leading cause of death in developed countries with a heavy load on people in developing countries by causing morbidity and mortality [12]. The emergence of various drug-resistant organisms limited therapeutic efficacy of many of currently available drugs which is partly attributable to the limitation in both number and structural variety of anti-bacterial drugs in clinical use. Thus, there should be a focus to be given to search bioactive molecules from plants, which could serve as hit or lead in antimicrobial drug discovery. *Dodonaea angustifolia* is one of the *Dodonaea* species that exist in Ethiopia, which is traditionally used for the treatment of various infectious diseases. It has been commonly used for the treatment of various ailments. The leave juice used as wound healing by the traditional healer. The fruit is crushed and mixed with honey and taken for a remedy against malaria and the steam bark is used for teeth brushing. However, there is no scientific

data pertaining to the phytochemical isolation of the natural product and pharmacological activities of the plant. Thus, the present work has focused on isolation, characterization and the evaluation of the bioactive molecules for their antibacterial activity from the leaves of the *Dodonaea angustifolia*

1.2. Objectives of the Study

1.3.1. General Objective

The main objective of this study was to identify secondary metabolites from the leaves of *Dodonaea angustifolia* and evaluate for antibacterial activities.

1.3.2. Specific Objectives

1. To isolate secondary metabolites from the leaves extract of *Dodonaea angustifolia* using chromatographic techniques;
2. To elucidate the structures of the isolated compounds using spectroscopic techniques, 1D and 2D NMR and,
3. To evaluate the antibacterial activities of the crude extracts and isolated compounds against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Enterococcus faecalis* bacterial strains.

1.4 Significance of the Study

The emergence and potential spread of resistance to currently available antibiotics, drugs, including penicillin and that all attempts to use high-throughput screening and combinatorial chemistry, genomics, and vaccine development have so far not delivered a single drug despite the enormous resources these fields received during the past 16 years. At present, there are no promising candidates in the pipeline, signifying the importance of searching for new lead structures in antimicrobial drug development.

In Africa, the use of Medicinal plants still plays an important role in microbial treatment. Taking into account that plants were potential sources of the existing first line antimicrobial drugs, there is still great potential of identifying antimicrobial drugs from plant-based sources. Thus, screening and identifying the bioactive constituents of traditional medicinal

plants empirically used to treat infectious diseases will have a great contribution in addressing this global problem. Flavonoids, could exhibited promising activity against different types of ailment.

Members of the subfamily Asapindaceae including the genus *Dodonaea* produce a broad variety of Flavonoids and isoflavones, with wide range of bioactivities. Thus, the study of *Dodonaea angustifolia* may yield promising antibacterial agents. Furthermore, the findings of this research could be useful to provide information of the chemical profile of the plant and can be used as a database and guideline for further isolation and purification of the active principles

2. Review of Related Literature

2.1. The Microbial Burden

Infectious diseases are the second leading cause of death worldwide, and the third leading cause of death in developed countries in both children and adults. The burden of infectious diseases falls most heavily on people in developing countries by causing morbidity and mortality [12]. The challenge associated with fighting these diseases has become an increasingly complex one, because of the fast development of resistance to the classic antibiotics; the emergency of multiple drug resistant human pathogenic microorganisms and the changing nature of the infections observed in the elderly and other immune-compromised patients. Antimicrobial resistance is natural biological phenomenon recognized first as a scientific curiosity and then as a threat to the effective treatment of most microbial infections [13]. This demands the search for new sources of antimicrobial substances mainly from medicinal plants, because plants constitute a potentially useful resource for new and safe drugs for the treatment of microbial infections and other diseases [14].

So, to prevent and control infectious diseases it is a vital interest and concern in both developed and developing countries to come up with new remedies for such diseases [13].

To overcome this, new antimicrobial agents with different modes of action against bacterial, fungal, parasite and viral diseases are urgently needed. Thus, screening of medicinal plants for biological activity has resulted in the isolation of active compounds from the early 19th century when morphine (**6**) was isolated from *Papaver somniferum*. The success of the isolation of morphine led to the identification of many other drugs such as quinine (**1**) from *Cinchona spp*, atropine (**7**) from *Atropa belladonna*, reserpine (**8**) from *Rauwolfi serpentina* and digoxin (**9**) from *Digitalis* species [12].




Figure 2. structure some plant based compound for antibacterial

2.1.1 Economical Burden of Infection Disease

Even though the burden of infection diseases has been steadily decreasing in past decades in developed countries, it is still considerable worldwide. Lower respiratory infections, diarrhoeal and HIV/acquired immunodeficiency syndrome (AIDS) diseases are still among the top major killers in 2011 and infection diseases in general are responsible for considerable morbidity in all parts of the world [15].

In 2011, the global prevalence of HIV accounted for 34 million people. 69 % of them lived in Sub-Saharan Africa. Around five million people are living with HIV in South-East and East Asia combined. Other high-prevalence regions include the Caribbean, Eastern Europe and Central Asia [16]. Worldwide, HIV incidence is in downturn. In 2011, 2.5 million people acquired HIV infection; this number was 20 % lower than in 2001. Sharpest declines in the incidence have been recorded in the Caribbean (42 %) and Sub-Saharan Africa (25 %). However, variation among regions gives rise to concerns; since 2001, a 35 % increase of HIV incidence has been reported in the Middle East and North Africa. The number of newly infected people in Eastern Europe and Central Asia has been scaling up since 2001, as well [16].

World wide, an estimated 499 million new cases of curable sexual transmitted infection (as gonorrhoea, chlamydia and syphilis) occurred in 2008; these findings suggested no improvement compared to the 448 million cases occurred in 2005. However, wide variations in the incidence of sexual transmitted infections are reported among different regions; the burden of STIs mainly occurs in low-income countries [17]. According to the crucial burden of chlamydia infection in Europe, the economic impact of chlamydia infection has been deeply investigated; in the UK, the cost of chlamydia complications has been estimated to a minimum of € 110 million, annually [18]. Each year, in the USA, direct costs of chlamydia and its complications range between € 1 and 3 billion.

With regard to costs of influenza, results of a 2007 study, referring to 2003 data, highlighted the huge economic brunt of the burden of influenza in the USA, accounting for US\$ 87.1 billion across all age groups [19].

It is estimated that in 2010 alone, malaria caused 216 million clinical episodes and 655,000 deaths. An estimated 91 % of deaths in 2010 were in the African Region, followed by 6 % in the South-East Asian Region and 3 % in the Eastern Mediterranean Region . About 86 % of deaths globally were in children. A total of 3.3 billion people (half the world's population) live in areas at risk of malaria transmission in 106 countries and territories [20, 21]. Malaria imposes substantial costs to both individuals and governments. Direct costs for malaria have been estimated to be at least US\$ 12 billion per year worldwide [20,21].

Diarrhoeal diseases cause for 4.1 % of the total disability-adjusted life years global burden of disease, and are responsible for 1.8 million deaths every year. An estimated 88 % of that burden is attributable to unsafe supply of water, sanitation and hygiene [22]. Children in the developing world are the most affected by diarrhoeal disease: It is estimated that diarrhoeal diseases account for one in nine child deaths worldwide, making diarrhoea the second leading cause of death among children under the age of 5 after pneumonia [23].

2.1.2 Bacterial Infections and Antibacterial Agent

Bacteria are widely distributed and the most abundant group of unicellular organisms on earth, capable of adapting in a diverse range of environments such as in soil, water and air. They are both useful and harmful to humans. Many parasitic bacteria do not harm their hosts. Some cause disease through different mechanisms including production of toxins and or hydrolytic enzymes. Bacteria have a prokaryotic cell type with a rigid wall which protects the cell against osmotic damage. The structure of the cell wall differs in Gram-positive and Gram-negative bacteria [24].

Gram-negative bacteria differ from gram-positive by the presence of an outer membrane composed of lipopolysaccharides. It also contains specific proteins for transporting hydrophilic molecules. Other proteins are receptor sites for phages and bacteriocins. The outer membrane in Gram-negative bacteria covers the peptidoglycan layer, which is attached to the outer membrane by lipoproteins. The layer is separated by periplasm from the cytoplasmic membrane. Gram-positive bacteria have thicker peptidoglycan layers with no periplasm[24]. *Staphylococcus aureus* is a spherical Gram-positive parasitic bacterium that

causes illnesses ranging from minor skin infections and abscesses, to life-threatening diseases such as pneumonia, meningitis and septicemia[26].

Escherichia coli is a Gram-negative bacterium normally present in the intestinal tract of humans and other animals. *Escherichia coli* can sometimes be pathogenic, thus posing a threat to food safety, causing diarrhea, wound and urinary infections [24].

The other problem associated with the emergence of various resistant organisms which limited therapeutic efficacy of many of the available drugs. In addition, it has a serious global economic impact that affects the productivity of individuals, families and the society at large, since it causes energy loss, debilitation and loss of work capacity. Consequently, there is an urgent need for new, affordable and accessible antibacterial agents from nature with novel mechanism of action. The emergence and potential spread of strains of bacteria which are resistant to currently available drugs, has actually prompted the search for new drugs through the use of high-through put screening and combinatorial chemistry, genomics, and vaccine development. However, this effort has yet to deliver a single drug despite the enormous resources expended during the past many years. Natural products will continue to serve as lead structures for the development of antibacterial drugs. This is in fact a pointer that new antibacterial leads may emerge from plants, especially from those with recognized traditional uses.

2.2. Botanical Information

The family Sapindaceae, also called soapberry family is a flowering plants consisting of about 2000 species in *ca* 150 genera that are mainly distributed in temperate and tropical regions of the world [27]. The majority of these plants are either shrubs or trees, which are widely used for the treatment of various ailments [28] including its uses as medication to reduce fever, stimulating metabolism uses as antioxidant, analgesic, antiviral, anti-inflammatory, antiulcer and antibacterial activities [29, 30]. It is described that around 10% of the genera including *Dodonaea* and 1% of the species under the Sapindaceae family were registered in Ethiopian flora [27]. *Dodonaea* is considered to be largest genus with 70 species most of which are limited to Australia, one in New Guinea *Dodonaea polyandra*, *Dodonaea*

madagascariensis endemic to Madagascar and *Dodonaea angustifolia* is widely distributed in almost all continents [31, 32].

2.2.1 *Dodonaea angustifolia*

Dodonaea angustifolia, hop-bush in English, *Kitkita* in Amharic and *Ittacha* in Afaan Oromo is mainly found in Australia, Africa, Asia and South America [33, 34] with its greatest center of diversity appears to be the Southeast Asian region. *Dodonaea angustifolia* is a multi stemmed shrub or single-stemmed small tree up to 7 m tall. It occurs in different African countries like Ethiopia, Kenya, Somalia, Senegal, Nigeria, Mozambique, Madagascar, Democratic Republic of Congo, and South Africa in natural habitat and home cultivated in Ghana, Nigeria and Cameroon. It is an evading plant in East Africa which reproduces itself from seed very freely and grows on dry rocky slopes between 1500 and 2100 m. [35]



Figure 3. Picture of *D. angustifolia* on uncultivated area taken from Estern Wollega ,Gobu Sayo woreda, Anno district in 2016.

2.2.2 Traditional Uses of the Genus *Dodonaea*

Some plants under the genus *Dodonaea* are known to be used for the treatment of a wide range of diseases and illness. The aerial parts of *Dodonaea angustifolia* are used in folk medicine against fevers, swellings and colds in Latin America, China, Africa, and India [36, 37] while the juice made from the powdered leave is used for the treatment of trachoma and to expel roundworms as anti-helmentic [30]. There is also a report indicating that various parts of this plant are used in the traditional systems of medicine for contraceptive effect [38]. In the Tigray region of Ethiopia, the crushed fruit mixed with honey and taken for a remedy against malaria [39]. The stems are also used as fumigants to treat rheumatism [30].


2.2.3 Compounds from Genus *Dodonaea*

Dodonaea has been reported to produce diverse groups of secondary metabolites including flavonoids, terpenoids, saponins, tannins, cardiac glycosides, steroids, etc [40]. The major secondary metabolites so far reported are flavonoids and terpenoids [41] are summarized in Table 1.


Table 1: List of compounds isolated from *Dodonaea* species.

Compound class	Compound name	Plant source	Ref
Flavonoids	3,5,7-Trihydroxy-4'-methoxyflavone (10)	<i>Dodonaea viscosa</i>	[42]
	5,7,4'-Trihydroxy-3,6-dimethoxyflavone(11)	“	[43, 44]
	5,7-Dihydroxy-3,6,4'-trimethoxyflavone/santin(12)	“	[43, 45]
	5-Hydroxy-3,7,4'-trimethoxyflavone (13)	“	[30]
	5-Hydroxy-3,6,7,4'-tetramethoxyflavone(14)	“	
	3,5-Dihydroxy-7,4',-dimethoxyflavone (15)	“	
	3,4',5,7-Tetrahydroxyflavone(16)	“	[43, 44]
	Pinocembrin (17)	“	[44]
Methylenebissantin (18)	“	[46]	


	Kaempferol-3-methylether (19)	“	[46]
	5,7,4'-Trihydroxy-3'-(3-methylbut-2-enyl)-3-methoxyflavones(20),	Dodonaea polyandra	[47]
	7-Dihydroxy-3'-(3-methylbut-2-enyl)-3,4'-dimethoxyflavone(21)	“	
	5,7,4'-Trihydroxy-3',5'-(3-methylbut-2-enyl)-3-methoxyflavone (22)	“	
	5,7,4'-Trihydroxy-3',5'-(3-methylbut-2-enyl)-3-methoxy-6methoxyflavone(23)	“	
	5,7,4'-Trihydroxy-3'-(3-methylbut-2-enyl)-3-methoxy-5'methoxyflavone (24)	“	
Phenolics	Vanillicacid (25)	Dodonaea.viscosa	[46]
	NebrodensideA (26)	“	
	Docosylcaffeate (27)	“	
Terpenoids	5-(2-Furan-3-yl-ethyl)-8 α -hydroxymethyl-5,6-dimethyl-3,4,4 α ,5,6,7,8,8 α -octahydro-naphthalene-1-carboxylicacid		[46]
	hautriwaiclactone (28)		[48]



11



(12)



13




Figure 4. Structures compound isolated from *Dodonaea* species

2.2.4. Biological Activities of the Genus *Dodonaea*

The alcohol and aqueous extracts of the root of genus *Dodonaea* significantly reduced diarrhea in mice with reduction in weight of stools showing its anti-diarrheal property [49]. The leaves extract of genus *Dodonaea* exhibited anti-diabetic [36, 42.50.51], anti-inflammatory [52, 53], antioxidant [31, 43, 45], hypotensive [30], antiviral [54], analgesic [30], antipyretic [55] and wound healing activities [56]. Furthermore, the crude extract of the seeds of *Dodonaea* showed anti-malarial activity [57]. Literature report showed that different solvent extracts of the various parts of *D. angustifolia* were capable of suppressing the germination and growth of various plants demonstrating its allelopathic potential [58].

3 Materials and Methods

3.1 Chemical, Apparatus and Equipment

Rotary evaporator (Heidolph, USA) for solvent evaporation, Uv chamber for detection of spots on TLC plate, glass column chromatography (300 mm (B-14/23,B-19/26) and 500 mm, B-34/35); Smith Scientific Eden bridge; U.K) for separation and purification techniques, mortar and pestle for grinding, and round bottom flask of size 250, 500, and 1000 mL, measuring cylinder, filter papers (cotton swab), weighting balances, oven for drying purpose, TLC analyses were carried out on Merck pre-coated silica gel 60 F₂₅₄ plates. Column chromatography was run on silica gel 60 Å (70-230 mesh). Chloroform (99%), methanol (99.99%), from Blulux Laboratories Ltd, petroleum ether and ethyl acetate (99%), from Fisher Scientific, UK are chemicals used for extraction and gradient column elution. Dimethyl sulfoxide (DMSO), Mueller Hinton agar and nutrient broth as culture media were used for antibacterial activity test. Deuterated chloroform (CDCl₃, 99.8%) and dimethylsulfoxide (DMSO, 99.96%) solvents were used for obtaining NMR spectra, Vertical Laminar Flow Cabinet hood CLB-201-14 and Genlab incubator were used in the study. NMR spectra were obtained on Bruker Avance III 400 spectrometer, using the residual solvent peaks as reference. The spectra were processed using MestReNova 11.0 software.

3.2 Collection and Preparation of Plant specimens

The leaves of *Dodonaea angustifolia* were collected from Eastern Wollega Zone, Gubu Sayo woreda, Anno district in April 2016. The plant material was air-dried under shade and powdered to suitable size by using mechanical grinder to improve the subsequent extraction by rendering the sample more homogenous, increasing the surface area, and facilitating the penetration of solvent into the cells.

3.3 Extraction and Isolation

3.3.1 Extraction of Leaves of *Dodonaea angustifolia*

The air-dried and powdered leaves of *Dodonaea angustifolia* (0.5 kg) was soaked in 1:1 (2.5 L each) ratio of (chloroform /methanol) and exhaustively extracted 72 hr. each 3x24 hr. The filtrate was then separated from the mark using fresh cotton plug and the mark was macerated three times with the same solvent system. The filtrate was concentrated using rotary evaporator at 50 °C under reduced pressure to yield 110 g (22 %,) black crude extract and stored at room temperature. The portion of the crude extract (90 g) was subjected to differential extraction between ethyl acetate and water in 8:2 ratios. The process (partitioning) was repeated three times using the same ratio of ethyl acetate and water (8:2) followed by concentration of the ethyl acetate layer. The ethyl acetate extracts were then combined and concentrated to afford the dried weight of 36 g from plant material and (40%) from crude extract.

3.3.2 Isolation of Leaves of *Dodonaea angustifolia* extract.

A 30 g portion of the ethyl acetate extract was absorbed on 6 g silica gel (60-130 mesh) and subjected to column chromatography (500 mm diameter) on silica gel (145 g) eluting with petroleum ether with increasing amounts of ethyl acetate gradient. The following ratio of solvent combination were used in the elution process in petroleum ether to ratio of 100:00, 99:1, 98:2, 96:4, 90:10, 85:15, 80:20, 70:30, 50:50, 30:70, 20:80, 00:100 for each ratio fifty fraction with 30 mL was collected.


After TLC analysis, the fractions from SF18-24 (3.4 g) collected with 20% ethyl acetate in petroleum ether was subjected to column chromatography again on silica gel using increasing amounts of ethyl acetate in petroleum ether to give 250 mg of compound (**1a**)

0.5 K g *Dodonaea angustifolia* material

- Collected and chopped
- Air dried and powdered (500 g)
- Soaked with Chloroform/Methanol (1:1; 2x2.5 L) for 3x24 hr.

110 g crude extract

↓ 90 g suspended on 80% ethyl acetate in water



3.4 Culture Media

3.4.1. Test Strains

Once the isolate bacterial strains were obtained, the antibacterial susceptibility test was done in the order of selection of the strain, prepare and standardize inoculum suspension, plate inoculated, antibacterial disks added, incubated the plates and measured the zone of inhibition for interpretation was considered to know the potency of plant extract.

In vitro antibacterial activity was examined for crude the two extracts and for the isolated compound 34 from leave of *D. angustifolia*. Among microorganisms investigated, the two gram-positive and two-gram negative bacterial strains, *Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 respectively.

The standard strains were obtained from, Medical Microbiology Research Laboratory, Jimma hospital, denoted previously from Ethiopia Public Health Institute, Addis Ababa, Ethiopia. The bacteria strains were reactivated by sub culturing in nutrient broth at 37 °C and maintained on nutrient agar slant at 4 °C for further use in microbiology laboratory, Department of Biology Jimma University.

3.4.2 Standardization of Inoculums

To standardize the inoculums density for a susceptibility test, a Barium sulphate (BaSO_4) turbidity standard, equivalent to a 0.5 McFarland standard, should be used. Accordingly, BaSO_4 0.5 McFarland standard was prepared, by adding 0.5 mL of 0.048 molL^{-1} barium chloride (BaCl_2) 1.175% w/v dihydrated barium chloride ($\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$) in to 99.5 mL of 0.18 mol/L Sulphuric acid (H_2SO_4 , 1% v/v) with constant stirring to maintain a suspension. This mixture was considered to be equivalent to cell density of 1 to 2×10^8 CFU/ml. The turbidity standard is then aliquot into test tubes identical to those used to prepare the inoculum suspension. McFarland turbidity standard tubes were sealed with para film to prevent evaporation. Barium sulphate turbidity was compared with identical tubes containing inoculum 0.85% NaCl saline solution [59]. For each bacterial strain, Gentamycin was taken as positive control and pure solvent (DMSO) as the negative control [60].

3.5 Antibacterial Activity Test

3.5.1 Agar Disc Diffusion Method.

Agar disc diffusion method was used to evaluate the antibacterial activities of the crude extract and isolated compound on nutrient agar. Accordingly, an overnight culture of the test strains whose cell densities were adjusted to 0.5 McFarland standards (equivalent to 1 to 2×10^8 CFU mL⁻¹ bacterial suspensions) were uniformly spread on pre-prepared sterile Muller Hinton Agar plates to form lawn cultures. Then, 0.2 g crude extracts of (1:1) chloroform/methanol were dissolved in DMSO. The stock solution was prepared with concentrations of 200 mg mL⁻¹ for each solvent extracts. The blotting paper discs (6 mm) was soaked in various dilute solvent extracts and dried for 5 min. to avoid flow of extracts in the test media. The discs were placed on a plate with standardized culture suspension at equidistance to one another to avoid overlap of zones of growth inhibitions. Antibacterial activity of potential plant extract against bacterial pathogens was determined after incubation of the plates for 24 hr. at 37 °C, by measuring the diameter of zone of inhibition of growth. The diameter of zone of growth inhibition surrounding the discs was measured and expressed in millimeter (mm) using transparent ruler [61]. Each experiment was carried out in triplicates. The mean of the inhibition zone of each test sample was taken for evaluating the antibacterial activity of the extract.

4. Results and Discussion

4.1 Secondary Metabolites from leaves of *Dodonaea angustifolia*

The air dried leaves of *Dodonaea angustifolia* was exhaustively extracted with 1:1 (CH₃OH/CHCl₃) by maceration at room temperature and gave 110 g (20%) yield of initial plant material (0.5 kg). A portion of crude extract was subjected to liquid-liquid partitioning between water and ethyl acetate yielded 36 g ethyl acetate extract. A portion of the ethyl acetate extract (30 g) was subjected to chromatographic separation on silica gel eluting with increasing gradient of ethyl acetate on petroleum ether. Chromatographic separation of the extract resulted in the isolation of 5,7-dihydroxy 3,6,4'-trimethoxyflavone(1a). The structures of these compound was determined by 1D and 2D (COSY, HSQC, HMBC) NMR and compared with reported literature.

4.2 Characterization of Isolated Compound (1A)

Compound **1A** was isolated as a yellow solid from the ethyl acetate soluble portion of the 1:1 chloroform/methanol extract from leaves of *D. angustifolia*. The ¹H NMR spectrum revealed the presence of three proton singlet signals at δ_{H} 3.84, 3.89, and 4.03, each integrated for three protons for three methoxy groups. It also further showed the presence of five aromatic proton signals at δ_{H} 6.54 (1H, s), δ 7.02 (2H, d, J = 7.01Hz) and δ 8.07 (2H, d, J = 8.66 Hz) and two hydroxyl proton signal at δ_{H} 12.91 (1H, s) and δ_{H} 6.66(1H, s).

The ¹³C-NMR spectrum revealed that ten quaternary carbons which are attributed to five oxygenated aromatic carbons at δ_{C} 151.8 (C-5), 130.1 (C-6), 152.2 (C-7), 155.1 (C-9) and 161.1 (C-4'); two oxygenated olefinic carbons at δ_{C} 138.5 (C-3) and 156.2 (C-2); two aromatic quaternary carbons at δ_{C} 106.2 (C-10) and 122.9 (C-1'); and an α , β -unsaturated carbonyl carbon at δ_{C} 179.3 (C-4) and five methine carbon signals are also observed in the aromatic region at δ_{C} 93.2 (C-8), 114.2 (C-3', 5') and 130.3 (C-2', 6'). Furthermore, the spectrum demonstrated the presence of three methoxy groups at δ_{C} 55.6, 60.2 and 61.0 as showed in Table 2 (appendix 3, 4)

Table 2: The ^1H and ^{13}C NMR spectral data of compound (**1A**) and literature for santin (**43**)

Position	Compound (1A)		Santin (43)		Remark
	δ_{H}	δ_{C}	δ_{H}	δ_{C}	
2		156.2		156.2	Quaternary
3		138.5		138.4	Quaternary
4		179.3		179.2	Quaternary
5		151.9		155.1	Quaternary
6		130.1		130.1	Quaternary
7		152.3		151.8	Quaternary
8	6.54 (1H,s)	93.2	5.57 (1H,s)	93.1	CH
9		155.1		152.2	Quaternary
10		106.3		106.1	Quaternary
1'		122.9		122.7	Quaternary
2'and 6'	7.02 (2H,d J=7.01)	130.3	7.02 (2H,d)	130.2	CH
3'and 5'	8.07 (2H,d, J=8.66)	114.2	8.07 (2H,d)	114.2	CH
4'		161.1		161.7	Quaternary
4'-OMe	3.89 (3H,s)	55.6	3.70 (3H,s)	55.5	CH ₃
3-OMe	3.84 (3H,s)	60.3	3.80 (3H,s)	60.2	CH ₃
6-OMe	4.03 (3H,s)	61.0	3.80 (3H,s)	60.9	CH ₃

Assessments made on the basis of COSY, HSQC and HMBC correlations; δ in ppm; Coupling constants are in Hz.

The 1H-1H COSY spectrum (Appendix 5) showed the correlation between two aromatic protons each symmetrically placed on Para substituted benzene ring H-2' at δ_{H} 7.01 and H-3' at δ_{H} 8.07. The two protons are found coupled to each other but no other 1H-1H correlations are observed from its COSY spectrum as illustrated in figure 5




Figure. 5 . Partial structure of compound (**1A**) based on Cosy spectrum

Whereas in ring A, only a single up field proton signal at δ_{H} 6.54 because of *diortho* oxygenation is characteristic for proton at C-8 (H-8), otherwise fully substituted with two hydroxyl and a methoxyl groups.

The downfield shifted proton signal at δ_{H} 12.91 (1H,s) for chelated hydroxyl group involved in hydrogen bonding was placed at C-5, *peri* to carbonyl group.

Table 3. HSQC data for compound **1A**.


Carbon (^{13}C NMR)		Proton(^1H NMR)		Multiplicity	Remark
No	δ_{C} (ppm)	No	δ_{H} (ppm)		
OMe-4'	55.6	H-OMe	3.89	S	CH ₃
OMe-3	60.3	H-OMe	3.84	S	CH ₃
OMe-6	61.0	H-OMe	4.03	S	CH ₃
8	93.2	8	6.54	S	CH
2'and 6'	130.3	2'and 6'	8.07	d (J = 8.66)	CH
3'and 5'	114.2	3'and 5'	7.02	d (J = 7.01)	CH

The HSQC spectrum showed correlation such that the methyl proton signals at δ 3.89, 3.84 and 4.03 correlate with the carbons at δ_{C} 55.6, 60.3 and 61.0, respectively and other correlations observed were between aromatic protons at δ_{H} 6.54, 7.02 (2H, d) and 8.07(2H, d) with aromatic carbon signals at δ_{C} 93.3, 114.2 (C-3' & c-5') and 130.3 (C-2' & c-6'), respectively as shown in table 3 (appendix 6)

Table 4. Observed correlations in HMBC data of compound 1A

Carbon	δ_C (ppm)	HMBC(1H - ^{13}C)
OMe-4'	55.6	H (4'-OMe') \leftrightarrow C-3', C-4'
OMe-3	60.3	H (3-OMe) \leftrightarrow C-3, C-4, C-10
OMe-6	61.0	H (6-OMe) \leftrightarrow C-5, C-6, C-7
C-8	93.2	H-3' \leftrightarrow C-7, H-8 \leftrightarrow C-9, H-8 \leftrightarrow C-10, H-8 \leftrightarrow C-6
C-3'	114.2	H-3' \leftrightarrow C-2', H-3' \leftrightarrow C-4', H-3' \leftrightarrow , H-3'
C-2'	130.3	H-2' \leftrightarrow C-3', H-2' \leftrightarrow C-1', H-2' \leftrightarrow C-2, H-2' \leftrightarrow C-6' H (5-OH) \leftrightarrow C-5, H (5-OH) \leftrightarrow C-6, H (5-OH) \leftrightarrow C-10
OH on C-5		
OH on C-7		H (7-OH) \leftrightarrow C-7, H (7-OH) \leftrightarrow C-6, H(7-OH) \leftrightarrow C-8,

The HMBC spectral analysis revealed that the proton signal at δ_H 6.54 (H-8) showed long range coupling with the carbon signals at δ_C 152.3 (C-7), 151.1 (C-9), 130.1 (C-6) and 106.3 (C-10). The most downfield shifted signal in the 1H NMR at δ_H 12.91 (5-OH) showed HMBC cross coupling with its neighboring carbons, C-5 (δ 151.9), C-6 (δ 130.1) and C-10 (δ 106.3) confirming the position of this hydroxyl group at C-5, as shown in figure 6 and appendix 7.

**Figure 6.** Partial structure of compound 1A based on HMBC spectra

The signal at δ_H 6.66 in 1H NMR spectrum showed no HSQC correlation indicating that the group is for phenolic hydroxylic group. It also shows that HMBC correlations with C-7, C-6 and C-8, supported the placement of the group at C-7 as indicated in figure 7.




Figure 7. Partial structure of compound **1A** based on HMBC spectra

The spectrum also showed significant correlation led to the conclusion that the methoxy groups at δ_H 4.03 and 3.84 are clearly attached to C-6 (δ_C 130.1) and C-3 (δ_C 138.5), respectively as showed in figure 7. The signal due to the third methoxy group at δ_C 3.89 correlates with the carbon signal at δ_C 161.2 (C-4') establishing the location of the third methoxy group at C-4' as showed in figure 32.




Figure 8. Partial structure compound **1A** based on HMBC spectra

The HMBC Spectra on ring B also showed the correlation of H-2'(δ_C 8.07) with C-3'(130.3), C-1'(122.9), C-2 (156.2), (C-6'(130.3) and H-3'(7.02) with C-4' (161.1),C-2'(130.3)/ The overall HMBC structure was given as indicated in table 3and figure 9.




Figure 9. Partial structure of compound **1A** based on HMBC spectra.

Finally from the assignments made on the basis of ^1H NMR, ^{13}C NMR, COSY, HSQC and HMBC correlations and spectra the structure of the isolated compound was deduced as in figure 10.




Figure 10. Final Structure of Isolated compound.

The above physical and spectral data of compound **1A** are in close agreement with the data presented in the literature for santin (5,7-dihydroxy 3,6,4'-trimethoxyflavone) [43], which was previously isolated from the leaves of *Dodonaea viscosa* [53], *Parthenium hysterophorus* [63] and the aerial parts of *Tanacetum microphyllum* [64].

4.2.1. Summary of Spectra and Physical data the Isolated Compound (1A).

Compound (**1A**): Yellow solid; soluble in methanol; TLC: R_f 0.64 (mobile phase EtOAc :petrol, (2:8); have molecular formula C₁₈H₁₆O₇; ^1H NMR (5000 MHz, CDCl₃): δ_{H} 3.84 (3H, s, OCH₃), 3.89 (3H, s, OCH₃), 4.03 (3H, s, OCH₃), 6.54 (1H, s, H-8), 7.02 (2H, d, $J = 8.80$, H-2' and 6'), 8.07 (2H, d, $J = 8.80$, H-3' and 5') and 12.91 (1H, s, 5-OH) and 6.66 (1H, s, 7-OH) and ^{13}C NMR (5000 MHz, CDCl₃) δ in ppm: C-2 (δ 156.2), C-3 (138.5), C-4 (179.3), C-5 (151.9), C-6 (130.1), C-7 (152.3) C-8 (93.2), C-9 (155.1), C-10 (106.3), C-1' (122.9), C-2', 6' (130.3), (C-3', 5') (114.2), C-4' (161.1), OCH₃ (61.0), OCH₃ (60.3) and OCH₃ (55.6).

4.3 Biological Assay

4.3.1 Antibacterial activities of the extracts and isolated compound of the leaves of *Dodonaea angustifolia*

The crude extracts from the leaves of *D. angustifolia* were evaluated against four pathogenic bacteria namely; *E. faecalis*, *S. aureus*, *E. coli* and *Ps. aeruginosa* using gentamycin as a positive control drug and DMSO as negative control (Table 5 and Appendix 1).

The zones of inhibition was measured and found to be fall in the range of NI-11 mm for Chloroform / methanol (1:1 v/v) extract, NI-22 mm for ethyl acetate soluble extract and 22-29 mm for gentamycin. Chloroform / methanol (1:1 v/v) extract did not inhibited the two gram positive and shows active inhibition zone on the gram negative bacterial strains. The ethyl acetate soluble extract showed inhibition zone on both gram positive and gram negative bacterial strains. The activities of ethyl acetate extract was observed to be comparable with the positive control drug against *S.aureus* strain and only half way active on *E. coli* when compared with gentamycin as illustrated in Table 5.

Table 5. Antibacterial activity of *Dodonaea angustifolia* leaves crude CHCl₃/CH₃OH, ethyl acetate soluble extracts and isolated compound.

Bacterial strains		Crude extract isolated compound and control zone of inhibition (mm).				
		CHCl ₃ /CH ₃ OH extract	EtOA-soluble partition.	Isolated compoud	G	DMSO
<i>E.faecalis</i>	+ve	NI	7	12	22	NI
<i>S.aureus</i>	+ve	NI	22	10	23	NI
<i>E.coli</i>	-ve	11	12	15	23	NI
<i>P.aeruginosa</i>	-Ve	9	NI	13	29	Ni

NI = No Inhibition, EtOA = ethyl acetate, G = gentamycin, DMSO= dimethyl sulfooxide, +ve = positive , -ve = Negative

The isolated compounds showed good activities against the tested bacterial strains. The diameter of growth inhibition zone displayed on both gram negative and gram positive were

good with variable degree of potency the tested compound. However, Compound (**1A**) has shown significant anti-bacterial activity on gram negative bacterial strains (*E. faecalis* and *Ps. aeruginosa*) than gram positive bacterial strains (*E. faecalis* and *S. aureus*). As compared to the standard positive control (Gentamycin), compound (**1A**) has displayed good activity on *E. coli* and weak activity on *S. aureus* comparatively.

5. Conclusion and Recommendation

5.1 Conclusion

It has been documented that *Dodonaea angustifolia* is widely used around the world as analgesic, anti-inflammatory, antiviral, antifungal, anti-ulcerogenic, spasmolytic, laxative, antimicrobial, hypotensive, rheumatism, gout, hemorrhoids, fractures and snake bites. In the course of this project work, one compound, **1A** was identified from the leaves of *Dodonaea angustifolia*. It was isolated from (8:2) petroleum ether/Ethyl acetate extract. The antibacterial activities of these compounds had effect on tested bacteria as compared to gentamycin hence accounting for the traditional use of this plant against bacteria. Moreover, the structural elucidation of this compound was done by using spectroscopic methods NMR and UV-Vis.

Therefore, the *in vitro* antibacterial activity tests revealed that antibacterial activity of the isolated compound from the leaves of *Dodonaea angustifolia* to be the promising source of bioactive compounds that could be used as lead compounds in search for new clinically effective antimicrobial compounds.

The characterized compounds was identified as 5,7-dihydroxy-3,4,6-trimethylflavonoid, trivial name santin, their occurrence in the leaves of *Dodonaea* was confirmed from the biosynthetic pathway.

5.2 Recommendation

- I. The phytochemical investigation of *D. angustifolia* yielded for the isolation of compounds from the genus further phytochemical investigation on *D. angustifolia* should be done using HPLC so as to check the presence of phenyllanthraquinones, which serves as a marker compound in the genus *Dodonaea*.
- II. The crude extracts and isolated compounds showed considerable clear zone inhibition on four bacteria strains *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, and *Pseudomonas aeruginosa*, the other phytochemicals from the genus should be done on another type of bacteria strain for better activity.
- III. The clear zone of inhibition measured for isolated compound on *E.coli* was 15 mm as compared to the standard gentamycin was about 23 mm in diameter, for this compound the minimum inhibition concentration should be done further.
- IV. The isolated compound showed more potency for gram-negative bacteria; in contrast to the ethyl acetate soluble extract was more active for gram positive bacteria rather than gram negative bacteria. thus, the mechanism of action of these compound should be studied further

5. Reference

1. Bako, S.P.; Bakfur, M.J.; John, I.; Bala, E. I. Ethnomedicinal and phytochemical profile of some savanna plant species in Nigeria. *Int. J. Bot.* **2005**, *2*, 147-150
2. WHO. Traditional Medicine. Media Centre Fact Sheet No 134. Accessed from <http://www.who.int/mediacentre/factsheets/fs134/en/> **2008**.
3. Barnes, P.M.; Bloom, B.; nahin, R.L. Complementary and alternative medicine use among adult and children: Unit states, **2007**.
4. Mahesh, B.; Satish, S. Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World. J. Agr. Sci.* **2008**, *4*, 839-843
5. Cragg, G. M.; Newman, D. J.; Snader. K. M. Natural products in drug discovery and development. *J. Nat. Prod.* **1997**, *60*, 52-60.
6. Craig, W. J. Health-promoting properties of common herbs. *Am. Clin. Nutr.* **1999**, *70*, p. 491S-499S,
7. Bouamama, H.; Noel, T.; Villard, J.; Benharref, A.; Jana, M. Antimicrobial activities of the leaf extract of two Moroccan *Cistus* L species. *J. Ethno.pharm.* **2006**, *104*, 104-107.
8. Arora D, S.; Kaur G,J. Antibacterial activity of some Indian medicinal plants. *J of Nat.Med.* **2007**, *61*, 313-317.
9. Yi-Fang, C.S.; Jie, W,U; Xian-Hong, Rui-Hai L. Antioxidant and Antiproferative activities of common vegetables. *Rev J Agric Food Chem* **2002**; *50*:6910-6916.
10. Yineger, H.; Yewhalaw, D. Traditional medicinal plant knowledge and use by local healers in Sekoru District, Jimma Zone, South western Ethiopia. *J. Ethnobiol. Ethnomed.* 2007, *3*, 1-7
11. Solomon, Ch.; Ugochukwu 1, Arukwe Uche I. Onuoha, I. Preliminary phytochemical screening of different solvent extracts of stem bark and roots of *Dennetia tripetala* G. Baker. *Asia. J. of Plant Scie. and Res.*, **3**, **2013**, :10-13
12. Hedgerg, I.; Edwards,S. Pittosporaceae to Araliaceae. Flora of Ethiopia, **1989**, *3*, 490–503

13. Pascaline, J.; Charles, M.; Lukhoba, C.; George, O., Phytochemical constituents of some medicinal plants used by the Nandis of South Nandi district, Kenya. *J An. Plant .Sci.* 9,1201-1210.
14. Teka, T.; Bisrat, D.; Mazumder, A.; Asres, K. Antimicrobial constituents from the leaf latex of aloe cherrima Gilbert & Sebsebe
15. The top 10 causes of death. Fact sheet N°310 (Updated July 2013) World Health Organization, Global burden of disease.
<http://www.who.int/mediacentre.fact.sheets.html>. Accessed 28 Jan 2015
16. UNAIDS report on the global AIDS epidemic. UNAIDS 2013.
17. WHO Sexually transmitted infections (STIs). World Health Organization, Geneva.2013
18. ECDC Chlamydia control in Europe. ECDC, Stockholm 2009
19. Molinari N.A.; Ortega-Sanchez, I.R.; Messonnier, M.L.; Thompson, W.W.; Wortley, P.M. The annual impact of seasonal influenza in the US: measuring disease burden and costs. *Vaccine* **2007**, 25, 27, 5086–5096
20. World Malaria Report 2011, WorldHealthOrganization.http://www.who.int/malaria/world_malaria_report_2011/en/index.html. Accessed 28 Jan 2015
21. The Global Malaria action Plan for a Malaria Free World. <http://www.rollbackmalaria.org/gmap/>. Accessed 28 Jan 2015 <http://www.rollbackmalaria.org/gmap/>. Accessed 28 Jan 2015
22. Cairncross, S.; Hunt. C.; Boisson, S.; Bostoen, K.; Curtis, V.; Fung, I.C. Schmidt WP Water, sanitation and hygiene for the prevention of diarrhoea. *Int. J. Epid.* **2010**, 39, 193– 205
23. Aiello AE, Coulborn RM, Perez V, Larson EL Effect of hand hygiene on infectious disease risk in the community setting: a meta-analysis. *Am J Public Health* **2008**, 98, 1372–1381
24. Sleight, J. D. Antibiotic resistance in *Serratia marcescens*. *Brit. Med. J (Clin res.)* **1983**,287 6406, 1651
25. Kumar, K.; Jha, S. Study Of Viability Period of Culture Suspension Of Salmonella Enterica.

26. Halablab, M.; Hijazi, S.; Fawzi, M.; Araj, G., Staphylococcus aureus nasal carriage rate and associated risk factors in individuals in the community. *Epid. and infection* 138 05, 702-706
27. Cook, A.; Turner, S.R.; Baskin, J.M.; Baskin, C.C.; Steadman, K.J.; Dixon, K.W. Occurrence of Physical Dormancy in Seeds of Australian Sapindaceae: A Survey of 14 Species in Nine Genera, *Annals of Botany*, **2008**. 101,1349–1362
28. Hamerski, L.; Somner, G.V.; Tamaio, N. Paullinia cupana (Sapindaceae): A Review of Its Ethnopharmacology, Phytochemistry, and Pharmacology. *J. Med. Plant. Res*, **2013**, 30, 2221–2229.
29. Rani, M.S.; Pippalla, R.S.; Mohan, P.K. Dodonaea viscosa Linn.-An Overview, *JPRHC*, **2009**, 1, 97–112.
30. Shepherd, K.A.; Rye, B.L.; Meissner, R.A.; West, J.G. Two New Western Australian Species of Dodonaea from Northern Yilgarn Ironstones. *Nuytsia*. **2007**, 17, 375– 384
31. Harrington, M.G.; Gadek, P.A. A species Well Travelled- The Dodonaea viscosa Complex Based on Phylogenetic Analysis of Nuclear Ribosomal ITS and TSF Sequences. *J. Biogeography*. **2009**, 36, 2313-2323.
32. Omosa, L.K.; Midiwo, J.O.; Derese, S.; Yenesew, A.; Peter, M.J.; Heydenreich, M. neo-15,16-Epoxy-2-hydroxy-3,13(16),14-clerodtriene-18-oic acids from the Leaf Exudate of Dodonaea angustifolia, *Phytochem.Lett.* **2010**.3,217–220.
33. Liu, J.; Noshiro, S. Lack of Latitudinal Trends in Wood Anatomy of D. viscosa A Species with World Wide Distribution. *Amer. J. Bot.* **2003**, 94, 532-539
34. Cunningham, P.L.; Joubert, D.F.; Adank, W. Dodonaea angustifolia an Alien Invasive to the Auas Mountains in Namibia. *Dinteria*. 2004, 29, 11–18
35. Al-Asmari, A.K.; Al Otaibi, K.E.; Al-Omani, S.; Athar, M.T. An Updated Phytopharmacological Review on Medicinal Plant of Saudi Arabia-Dodonaea viscosa Linn. *Am. J. Res. Comm.* **2013**, 12, 519–531
36. Naidoo, R.; Patel, M.; Gulube, Z.; Fenyvesi, F. Inhibitory Activity of Dodonaea viscosa Extract Against Streptococcus mutans and its Biofilm. *J. Ethn. phar.* **2012**, 144, 171–174
37. Ramya, R.; Sivasakthi, R.; Senthilkumar, S.; Anudeepa, J.; Santhi, N.; Narayanan, R.V. Preliminary Phytochemical and Anti-fertility Studies on Dodonea viscosa. *Asia. J. Res. Pharm. Sci.* **2011**, 3, 77–79


38. Yirga, G. Assessment of Indigenous Knowledge of Medicinal Plants in Central Zone of Tigray, Northern Ethiopia. *Afr. J. Plant. Sci.* **2010**, 4, 006–011
39. Pirzada, A.; Shaikh, W.; Usmanghani, K.; Mohiuddin, E. Antifungal Activity of *Dodonaea viscosa* Jacq Extract on Pathogenic Fungi Isolated from Super Facial Skin Infection, *Pak. J. Pharm. Sci.* **2010**, 23, 337–340.
40. Meenu, J.; Sunil, S.; Manoj, K. Evaluation of Antihyperglycemic Activity of *Dodonaea Viscosa* Leaves in Normal and Stzdiabetic Rats. *Int. J. Phar. Sci*, **2011**, 3, 69 –74
41. Prakash, N.K.U.; Selvi, C.R.; Sasikala, V.; Dhanalakshmi, S.; Prakash, S.B.U. Phytochemistry and Bio-Efficacy of a Weed, *Dodonaea viscosa*, *Int. J. Phar. Sci.* **2012**, 4, 509–512.
42. Veerapar, V.P.; Prabhakar, K.R.; Thippeswamy, B.S.; Pansal, P.; Sirinvasan, K.K.; Unikrshanan, M.K. Antidiabetic Effect of *D. viscosa* Aerial Parts in High Fructose Fedinsulin Resistant Rats mechanism Based Study, *Ind. J.Exp. Biol.* **2010**, 48, 800–810
43. Riaz, T.; Abbasi, M.A.; Rehman, A.A.; Shahzadi, T.; Ajaib, M.; Khan, K.M. Phytochemical Screening, Free Radical Scavenging, Antioxidant Activity and Phenolic Content of *Dodonaea viscosa* Jacq. *J.Serb.Chem.Soc.* **2012**, 77, 423–435
44. Muhammad, A.; Anis, I.; Ali, Z.; Awadelkarim, S.; Khan, A.; Khalid ,A.; Shah, M.R.; Galal, M.; Khan, I.A.; Choudhary, M.H. Methylenebissantin: A rare Methylene-Bridged Bisflavonoid from *Dodonaea viscosa* which Inhibits Plasmodium falciparum enoyl-ACP reductas. *Bioorganic & Medicinal Chemistry Letters.* 2012, 22,610–612
45. Oliveira, S.Q.; de Almeida, M.T.R.; Maraslis, F.; Silva, I.T.; Sincero, T.C.M.; Palermo, J.A.; Cabrera, C.M.; Caro, M.S.B.; Simoes, C.M.O.; Schenkel, E.P. Isolation of Three New ent-Labdane Diterpenes from *Dodonaea viscosa* (Sapindaceae): Preliminary evaluation of antiherpes activity, *Phytochemistry Letters*, 2012, 5,500–505.
46. Wabo, H.K.; Chabert, P.; Tane, P.; Noté,O.; Tala, M.F.; Peluso,J.; Muller, C.; Kikuchi, H.; Oshima, Y.; Lobstein, A. Labdane-type Diterpenes and Flavones from *Dodonaea viscosa*, *Fitoterapia*, 2012, 83, 859–863
47. Ortega, A.; Garcia, E.P.; Cardenas, C.; Marquina, S.; Garduno, L.D.; Maldonado, E. Methyl dodonates, a New Type of Diterpenes with Modified Clerodane Skeleton from *Dodonaea viscosa*, *Tetrahedron*, 2001.57,2981–2989

48. Asres, K.; Bucar, F.; Kartnig, T.; Witvrouw, M.; Pannecouque, C.; De Clercq, E. Antiviral activity against human immunodeficiency virus type 1 (HIV-1) and type 2 (HIV-2) of ethno botanically selected Ethiopian medicinal plants. *Phytoth. Res* 2001,15,62-69
49. Pirzada, A.; Shaikh, W.; Usmanghani, K.; Mohiuddin, E. Antifungal Activity of *Dodonaea viscosa* Jacq Extract on Pathogenic Fungi Isolated from Super Facial Skin Infection, *Pak. J. Phar.Sci.* **2010.**,23, 337–340
50. Akhtar, M.S.; Ahmed, M.; Gulzar, K.; Adnan, H. Hypoglycemic Activity of *Dodonaea viscosa* Leaves in Normal and Alloxan-Induced Diabetic Rabbits, *Diabetologia Croatica*, **2011**, 40, 71–80.
51. Khalil, N.M.; Sperotto, J.S.; Manfron, M.P. Anti-inflammatory Activity and Acute Toxicity of *Dodonaea viscosa*, *Fitoterapia*, **2006**, 77,478–480.
52. Getie, A.; Gebre-Mariam, T.; Rietz, R.; Huschka, C.; Schmidtke, M.; Abate, A.; Neubert, R.H.H. Evaluation of the Anti-microbial and Anti-inflammatory Activities of the Medicinal Plants *Dodonaea viscosa*, *Rumex nervosus* and *Rumex abyssinicus*, *Fitoterapia*, **2003**,74, 139–143.
53. Teffo, L.S.; Aderogba, M.A.; Eloff, G.N. Antibacterial and Antioxidant Activities of Four Kaempferol methyl ethers Isolated from *Dodonaea viscosa* var. *angustifolia* leaf extracts, *S. Afr.J.Botany*, **2009**, 1–5.
54. Anilreddy, B. Preparation, Characterization and Biological Evaluation of Some Overview of *Dodonaea viscosa*, *J. Phar. Sci. and Tech.*, **2009**, 1, 1–9.
55. Ramya, R.; Anudeepa, J.; Senthilkumar, C.; Rajendran, S.S.; Sivasakthi, R; Moorthy, C.; Venkatnarayanan, D.R. Wound Healing Activity of *Dodonea viscosa* Linn Ointment In Rats, *Int.J.Res.in Phar and Chemistry*, **2011**, 3, 481–484.
56. Mengiste, B.; Makonnen, E.; Urga, K. In vivo Antimalarial Activity of *Dodonaea angustifolia* Seed Extracts Against *Plasmodium berghei* Mice Model, *MEJS.*, 2012, 4, 47–63.
57. Rajeswari, D.T.; Gajalakshmi, S.; Jayanthi, P.; Vijayalakshmi. S.. Pharmacological Activities of *Dodonaea viscosa*, *J. Phar.Res*, **2011**, 10, 3331–3333
58. Vanheerden, F.R.; Viljoen, A.M.; van Wyk, B.E.. The Major Flavonoid of *Dodonaea angustifolia*, *Fitoterapia*, 2000, 71,602–604.


59. Riaz, T.; Abbasi, M.A.; Rehman, A.A.; Shahzadi, T.; Ajaib, M.; Khan, K.M.. Phytochemical Screening, Free Radical Scavenging, Antioxidant Activity and Phenolic Content of *Dodonaea viscosa* Jacq., *J.Serb.Chem.Soc.*, **2012**, 77,4,423–435
60. Oliveira, S.Q.; de Almeida, M.T.R.; Maraslis, F.; Silva, I.T., Sincero, T.C.M., Palermo, J.A., Cabrera, C.M., Caro, M.S.B., Simoes, C.M.O., Schenkel, E.P.. Isolation of Three New ent-Labdanone Diterpenes from *Dodonaea viscosa* (Sapindaceae): Preliminary evaluation of antiherpes activity, *Phytochemistry Letters*, 2012, 5, 500–505
61. Anilreddy, B. Preparation, Characterization and Biological Evaluation of Some Overview of *Dodonaea viscosa*, *J. of Phar. Sci. and Tech.*, **2009**,1, 1–9
62. Leonidah, K. Omosa, Beatrice Amugune, Beth Ndunda Trizah K. Milugo, Matthias Heydenreich, Abiy Yenesew, Jacob O. Midiwo. Antimicrobial flavonoids and diterpenoids from *Dodonaea angustifolia*. *S. Afr. J. Bot.* **2014**, 91, 58–62.
63. Roy, D.C.; Shaik, M.M. Toxicology, Phytochemistry, Bioactive Compounds and Pharmacology of *Parthenium hysterophorus*, *J. Med. Pla. Studies*, **2013**, 140, 126–141.
64. Martinez, J.; Silvan, A.M.; Abad, M.J.; Bermejo, P.; Villar, A. Isolation of Two Flavonoids from *Tanacetum microphyllum* as PMA-Induced Ear Edema Inhibitor, *J. Nat. Prod.* **1997**, 60, 142–144.

7. Appendices


Appendix 1. Bioassays Tests of Crude Extracts Zone of Growth Inhibition in mm




E.Feli



E.coli




S.aureus




P.aeruginosa


Appendix 2. Bioassays Tests of Isolated Compound Zone of Growth Inhibition in mm




E.Feli



E.coli




S.aureus




P.aeruginosa


Appendix 3. $^1\text{H-NMR}$ Spectrum of compound (1A) in CDCl_3




Appendix 4. $^{13}\text{C-NMR}$ Spectrum of Compound (1A) in CDCl_3




Appendix 5. COSY Spectrum of Compound (1A) in CDCl₃



Appendix 6. HSQC Spectrum of compound (**1A**) in CDCl₃





Appendix 7. HMBC Spectrum of compound (**1A**) in CDCl_3

