

## Full Length Research Paper

# Responses of fresh leaf yield and quality variables of $M_5$ mulberry to bio-inoculants, farm yard manure (FYM) and inorganic fertilizers under rain-fed conditions

Waktole Sori<sup>1\*</sup> and Bhaskar R. N<sup>2</sup>

<sup>1</sup>Department of Sericulture, University of Agricultural Sciences, GKVK, Bangalore-65- Karnataka-India.

<sup>2</sup>College of Sericulture, Chintamani-Karnataka, University of Agricultural Sciences, India.

Accepted 25 September, 2012

An experiment was conducted to know the effect of integrated use of bio-inoculants (*Azotobacter* sp. at 20 kg/ha/year, *Aspergillus awamori* at 25 kg/ha/year and *Trichoderma harzianum* at 20 kg/ha/year), farm yard manure (FYM) and inorganic fertilizers (IF, that is, Nitrogen-phosphorus-potassium (NPK) on fresh leaf yield and quality variables of  $M_5$  mulberry under rain-fed condition during 2006-07. The experiment was laid out using randomized complete block design with eight treatments replicated thrice. The results revealed that combined use of treatment components had a positive effect on fresh leaf yield and quality variables of mulberry on par with the standard check (100:50:50 kg/ha/year NPK and 12 MT/ha/year FYM). Application of the recommended dosage of fertilizers recorded maximum fresh leaf yield per plant (403.60 and 538.13; 718.74 and 867.57 g) at 45 and 60 days after pruning in the first and second crops, respectively. However, it was consistently and statistically on par with  $T_4$  (397.80 and 530.40; 614.07 and 795.20 g). Further, the leaf quality variables namely, moisture percentage (71.89 and 70.90; 71.50 and 68.90 at 45 and 60 days after pruning for crop I and II, respectively), chlorophyll (a) (1.45 mg/g), chlorophyll (b) (0.69 mg/g), total chlorophyll (2.14 mg/g), crude protein (18.21 mg/g), total soluble protein (8.82 mg/g), total soluble sugar (7.95 mg/g) and NPK (2.98, 1.72 and 1.33%, respectively) content of the leaf were maximum in the standard check which was on par with  $T_4$ . Thus, N and P inorganic fertilizers can be reduced to the tune of 25% without affecting the fresh leaf yield and quality variables of  $M_5$  mulberry by amending the same through bio-inoculants under rain-fed conditions.

**Key words:**  $M_5$  mulberry, bio-inoculants, yield and quality variables, rain-fed condition.

## INTRODUCTION

Mulberry (*Morus* spp.) is the sole food plant for the silkworm (*Bombyx mori* L.). Besides climate, rearing technique, silkworm race and silkworm eggs, are known to influence successful cocoon production by mulberry silkworm (Singhal et al., 1999). Thus, nutritive value of mulberry leaf plays a predominant role in better crop performance of mulberry silkworm. However, the nutritive value of mulberry leaf is influenced by the soil nutrient

status (Bongale, 2006). Frequent use of inorganic fertilizers for a prolonged period results in poor soil health and affects availability of nutrients to be absorbed by plants (Subbaswamy et al., 1994). It was reported that feeding of nutrient deficient mulberry leaves to developing silkworms led to 17.11% cocoon production loss (Singhvi et al., 2007). Use of 20 kg/ha/year of *Azotobacter* was effective in minimizing the application of

\*Corresponding author. E-mail: waktoletufa@yahoo.com.

**Table 1.** Month wise weather data in the study area during the study period.

Months	Total rain fall (mm)	Temperature (°C)		Relative humidity (%)		Wind speed (km/h)
		Maximum	Minimum	Morning	Noon	
<b>Year 2006</b>						
June	112.60	29.20	19.60	90.00	49.00	10.60
July	84.40	28.10	19.30	90.00	52.00	14.10
August	63.20	28.50	19.00	93.00	49.00	11.10
September	61.00	29.20	19.40	91.00	47.00	9.00
October	36.00	28.80	18.70	91.00	48.00	8.00
November	111.80	27.00	17.80	95.00	53.00	7.10
December	1.80	26.40	13.70	93.00	49.00	8.10
<b>Year 2007</b>						
January	0.00	28.10	13.80	90.00	42.00	7.70
February	0.80	29.80	14.80	90.00	40.00	7.90
March	0.00	33.20	18.40	82.00	33.00	7.60
April	109.60	33.80	20.20	83.00	33.00	7.50
May	85.40	33.20	21.20	85.00	38.00	9.10
June	60.30	29.60	19.70	91.00	49.00	11.20
July	149.20	28.40	19.90	90.00	50.00	12.10

inorganic fertilizers 'N', thereby, reducing the cost of mulberry cultivation under irrigated conditions without adverse effect on leaf quality (Das et al., 1994). Therefore, it was deemed necessary to test the influence of bio-inoculants on different functions of mulberry yield and quality variables under rain-fed conditions by complementing it with inorganic fertilizers and farm yard manure (FYM).

## MATERIALS AND METHODS

The experiment was superimposed on already established 10 years old *M<sub>5</sub>* mulberry to determine the influence of bio-inoculants (*Azotobacter* sp. at 20 kg/ha/year, *A. awamori* at 25 kg/ha/year and *T. hazianum* at 20 kg/ha year), FYM at 12 MT/ha/year and inorganic fertilizers (N from urea, P from single super phosphate and K from muriate of potash at 100:50:50 kg/ha/year and reduced dosages of NP plus the bio-inoculants) on fresh leaf yield and quality variables of *M<sub>5</sub>* mulberry under rain-fed condition at Gandhi Krishi Vignana Kendra. The study was established at Bangalore in the Department of Sericulture premises which is located at a coordinates of 77°35' E longitude and 12°58' N latitude at an altitude of 930 m above mean sea level. The weather data on rainfall, temperature, relative humidity and wind speed that prevailed during the cropping period were compiled and presented in Table 1. The rain fall was maximum in the month of July 2006/2007 (149.20 mm) and there was no rain during January and March 2006/2007. The temperature ranged between 13.70 to 33.80°C; the hottest month being April (33.80°C (Max.)) and 20.20°C (min.); however, the coldest month was December (26.40°C (Maximum) and 13.70°C (minimum)). Maximum relative humidity of 95.00% was recorded during November and minimum was during March (82.00%). Maximum wind speed of 14.10 km/hr was recorded in July but minimum of 7.10 km/h was registered in the month of November.

Bio-inoculants were obtained from bio-fertilizers scheme, Department of Agricultural Microbiology, UAS (B), GKVK and were mixed with FYM under wet condition and applied to the root zone of

mulberry two weeks after the application of inorganic fertilizers to avoid possible contact. The experiment was laid-out in a randomized complete block design with eight treatments each with three replications. A single plot size of 4 x 4 m (16 m<sup>2</sup>) was used for each replication. A spacing of 60 x 60 cm between plants and rows were maintained for the experiment. The treatments were: T<sub>1</sub> (recommended rate, that is, 100:50:50 kg/ha/year NPK and 12 MT/ha/year FYM), T<sub>2</sub> (75% of recommended NP from inorganic fertilizers, 25% of recommended NP from *A. awamori* and *Azotobacter* sp., 12 MT/ha/year FYM and 50 kg/ha/year K), T<sub>3</sub> (50% of recommended NP from inorganic fertilizers, 50% of recommended NP from *A. awamori* and *Azotobacter* sp., 12 MT/ha/year FYM and 50 kg/ha/year K), T<sub>4</sub> (75% of recommended NP from inorganic fertilizers; 25% of recommended NP from *A. awamori*, *Azotobacter* sp. and *T. hazianum*; 12 MT/ha/year FYM and 50 kg/ha/year K), T<sub>5</sub> (50% of recommended NP from inorganic fertilizers; 50% of recommended NP from *A. awamori*, *Azotobacter* sp. and *T. hazianum*; 12 MT/ha/year FYM and 50 kg/ha/year K), T<sub>6</sub> (recommended NPK from FYM), T<sub>7</sub> (Recommended NPK from inorganic fertilizers) and T<sub>8</sub> (Control, that is, no application of fertilizer).

Cultural practices were followed as per the package of practices for rain-fed mulberry as described by Dandin et al. (2003). Bottom pruning was carried out during the last week of June, 2006 and 2007 just at the onset of main rainy season. After one and two weeks of bottom pruning the inorganic fertilizers, bio-inoculants and FYM were applied as per the treatment details. Observations on fresh leaf yield and moisture percentage were recorded by harvesting the fresh leaves from five randomly selected and tagged plants from each plot 45 and 60 days after bottom pruning for two seasons. Leaves were harvested individually plant wise, weighed and then mean leaf yield per plant was determined. But observation on other quality variables viz., chlorophyll, crude protein, total soluble protein, total soluble sugars and NPK content of the leaf were made on the second crop 60 days after bottom pruning.

Leaf quality variables viz., moisture, chlorophyll (a), chlorophyll (b) and total chlorophyll, NPK, crude protein, total soluble proteins and total soluble sugar contents of the leaf were each estimated through analysis. Moisture percentage was determined by A.O.A.C. (1970), chlorophyll content by Hiscox and Israelstam (1979) and

**Table 2.** Fresh leaf yield responses of M<sub>5</sub> mulberry to integrated application of bio-inoculants, FYM and Inorganic fertilizers under rain-fed conditions

Treatments**	Fresh leaf yield/plant (g)			
	Crop I		Crop II	
	45DAP***	60DAP	45DAP	60DAP
T <sub>1</sub>	403.60	538.13	718.74	867.57
T <sub>2</sub>	397.00	529.33	526.00	605.70
T <sub>3</sub>	373.60	498.13	421.33	452.71
T <sub>4</sub>	397.80	530.40	614.07	795.20
T <sub>5</sub>	390.80	521.07	503.92	581.60
T <sub>6</sub>	360.20	480.27	364.20	497.10
T <sub>7</sub>	373.20	497.60	323.07	377.77
T <sub>8</sub>	257.00	342.67	226.77	280.05
F-test	*	*	*	*
SEm±	10.37	13.82	57.912	78.24
LSD <sub>5%</sub>	31.45	41.93	175.68	237.34
CV (%)	4.86	4.84	21.70	25.23

\*Significant at 5% level of significance; \*\*\*DAP-Days after pruning, \*\*T<sub>1</sub>-Standard check (RNPK + RFYM), T<sub>2</sub>- 75 % NP through IF and 25% through *A. awamori* and *Azotobacter* sp. + RFYM + RK T<sub>3</sub>- 50% NP through IF and 50% NP through *A. awamori* and *Azotobacter* sp. + RFYM + RK, T<sub>4</sub> - T<sub>2</sub> + *T. harzianum*, T<sub>5</sub>- T<sub>3</sub> + *T. harzianum*, T<sub>6</sub>- RNPK from FYM, T<sub>7</sub>- RNPK only from IF, T<sub>8</sub>-control.

Arnon (1949), total soluble protein was estimated as per the procedure of Lowry et al. (1951), crude protein content by multiplying the percentage of nitrogen in the leaf by a factor 6.25, total soluble sugar by colorimetric method (Dubios et al., 1956) and NPK percentage using standard procedures. Data were analyzed using ANOVA as described by Cochran and Cox (2000). Least significant difference (LSD) test were used to differentiate statistical mean differences at 5% level of significance whenever the means were found significant for pair wise comparison of the treatments.

## RESULTS AND DISCUSSION

### Fresh leaf yield per plant (g/plant)

The treatments were significantly ( $P < 0.05$ ) different from each other with respect to the fresh leaf yield of M<sub>5</sub> mulberry (Table 2). In the first and second crops, T<sub>1</sub> has recorded maximum fresh leaf yield per plant (403.60 and 538.13; 718.74 and 867.57 g/plant), which was consistently and statistically on par with T<sub>4</sub> (397.80 and 530.40; 614.07 and 795.20 g/plant) 45<sup>th</sup> and 60<sup>th</sup> days after pruning, respectively. Minimum fresh leaf yield (257.00 and 342.67; 226.77 and 280.05 g/plant) was recorded from the control 45<sup>th</sup> and 60<sup>th</sup> days after pruning (Table 1). The increased fresh leaf yield of M<sub>5</sub> mulberry from T<sub>1</sub> and T<sub>4</sub> may be due to more number of shoots, higher plant height and more number of leaves per plant. Further, the combined application of FYM might have helped in slow and steady release of nutrients in addition to supply of important macro- and micro-nutrients besides

efficient supply of N and P by nitrogen fixing and phosphorus solubilizing bio-inoculants utilized. Thus, NP application can be curtailed to the tune of 25% by supplementing with bio-inoculants without adverse effect on fresh leaf yield of M<sub>5</sub> mulberry under rain-fed condition. Katiyar et al. (1995) in their earlier studies on the effect of dual inoculation of *A. chroococum* and *G. mosseae* with 50% recommended dosage of N and P fertilizers have reported that, the yield of mulberry was on par with uninoculated control receiving full dosage of N and P fertilizers. Das et al. (1994) in their studies have also reported that, there was no reduction in leaf yield in K<sub>2</sub> variety of mulberry when 50% nitrogen fertilizer was compensated with 20 kg *Azotobacter* sp./ha/year. Sreeramulu et al. (2004) also reported a saving of 25% N and P chemical fertilizers application without affecting leaf yield of mulberry when *A. brasilense* and *B. megaterium* were inoculated.

### Moisture content (%)

The moisture content of M<sub>5</sub> mulberry leaves was significant ( $P < 0.05$ ) due to the influence of the different treatments in both the crops 45<sup>th</sup> and 60<sup>th</sup> days after pruning (Table 3). Among the different treatments, in both crops, the leaves harvested from T<sub>1</sub> plot recorded maximum moisture percentage (71.89 and 70.90; 71.50 and 68.90), respectively 45<sup>th</sup> and 60<sup>th</sup> days after pruning. This treatment was statistically similar with T<sub>4</sub> (71.08 and

**Table 3.** Moisture, chlorophyll and crude protein contents of M<sub>5</sub> mulberry as influenced by integrated application of bio-inoculants, FYM and inorganic fertilizers under rain-fed condition.

Treatments**	Moisture (%)				Chlorophyll (mg/g)		
	Crop I		Crop II		Chl.'a'	Chl.'b'	Total chl.
	45DAP***	60DAP	45DAP	60DAP			
T <sub>1</sub>	71.89	70.90	71.50	68.90	1.45	0.69	2.14
T <sub>2</sub>	71.00	69.96	70.00	67.84	1.42	0.64	2.06
T <sub>3</sub>	69.85	69.15	67.00	66.95	1.36	0.60	1.96
T <sub>4</sub>	71.08	70.13	70.75	68.13	1.44	0.68	2.13
T <sub>5</sub>	69.64	69.25	69.00	67.39	1.41	0.63	2.04
T <sub>6</sub>	67.99	67.09	68.00	66.26	1.36	0.63	1.99
T <sub>7</sub>	69.52	68.61	66.56	66.05	1.30	0.58	1.87
T <sub>8</sub>	65.56	64.92	62.56	62.82	1.24	0.48	1.71
F-test	*	*	*	*	*	*	*
SEm±	0.32	0.47	0.571	1.328	0.016	0.010	0.020
LSD <sub>5%</sub>	0.98	1.44	1.731	4.029	0.048	0.031	0.062
CV (%)	0.81	1.19	1.450	3.444	1.984	2.871	1.768

\*Significant at 5% level of significance; \*\*\*DAP-Days after pruning, \*\*T<sub>1</sub>. Standard check (RNPK + RFYM), T<sub>2</sub>. 75% NP through IF and 25% through *A. awamori* and *Azotobacter* sp. + RFYM + RK T<sub>3</sub> – 50% NP through IF and 50% NP through *A. awamori* and *Azotobacter* sp. + RFYM + RK, T<sub>4</sub> - T<sub>2</sub> + *T. harzianum*, T<sub>5</sub> – T<sub>3</sub> + *T. harzianum*, T<sub>6</sub> – RNPK from FYM, T<sub>7</sub> – RNPK only from IF, T<sub>8</sub> – control.

70.13%; 70.75 and 68.13%) and T<sub>2</sub> (71.00 and 69.96%; 70.00 and 67.84%) consistently. Minimum moisture content (65.56 and 64.92%; 62.56 and 62.82%) was registered from T<sub>8</sub>. Increased leaf moisture percentage may be due to the greater ability of the combined usage of the bio-inoculants, FYM and IF to support better plant performance by fixing more nitrogen and making available the bound phosphorous thereby improving water absorbing capacity of the roots which in turn has been translocated to the leaf. Better moisture percentage of mulberry leaf as a result of bio-inoculants application to mulberry plant was reported by Patil et al. (2001).

### Chlorophyll content (mg/g)

Maximum chlorophyll (a), (b) and total chlorophyll content of 1.45, 0.69 and 2.14 mg/g, respectively were recorded from leaves of mulberry plants that received T<sub>1</sub> which was followed and on par with T<sub>4</sub> (1.44, 0.68 and 2.13 mg/g) whereas minimum content of all the chlorophyll were registered from T<sub>8</sub> (1.24, 0.48 and 1.71 mg/g) after 60 days of pruning (Table 3). The positive influence of bio-inoculants, FYM and IF on the chlorophyll content was in conformity with the findings of Sunil (2005), who observed that, higher chlorophyll content in mulberry leaves was due to addition of organic manures, biofertilizers and inorganic fertilizer. The findings of Singhal et al. (2000), who also reported that application of nitrogen helped in harnessing more solar energy through chlorophyll synthesis corroborates the present findings. Shivakumar et al. (2000) revealed that the

organic manure (FYM) supplemented with varied levels of inorganic fertilizers caused increment in chlorophyll content of leaves in mulberry.

### Crude protein and total soluble protein content (mg/g)

The crude protein and total soluble protein content were significantly (P<0.05) more in T<sub>1</sub> (18.21 and 8.82 mg/g, respectively) 60 days after pruning (Table 4). The next best treatment was T<sub>4</sub> (17.71 and 8.47 mg/g). The lowest of the same were observed in T<sub>8</sub> (15.61 and 5.23 mg/g) (Tables 3 and 4). Katiyar et al. (1995) and Umesh (1999) reported similar leaf chemical constituent in mulberry when biofertilizers were inoculated with reduced chemical fertilizers. Curtailment of 50% N application without affecting mulberry leaf quality by supplementing mulberry through 20 kg of *Azotobacter* has been reported by Das et al. (1994).

### Total soluble sugar (mg/g)

Total soluble sugar was significantly (P<0.05) different among the different treatments (Table 4). The highest total soluble sugar content was found in the plot that received treatment T<sub>1</sub> (7.95 mg/g), which was statistically on par with T<sub>4</sub> (7.72 mg/g). The lowest total soluble sugar content of 5.32 mg/g was recorded in T<sub>8</sub>. Similar findings were reported earlier by many scientists, notably (Shankar et al., 2000; Das et al., 1994) stated quality variables of mulberry were not adversely affected when

**Table 4.** Total soluble protein, total soluble sugar and NPK contents of M<sub>5</sub> mulberry as influenced by integrated application of bio-inoculants, FYM and inorganic fertilizers under rain-fed conditions

Treatments**	Crude protein (mg/g)	Total soluble protein (mg/g)	Total soluble sugar (mg/g)	N (%)	P (%)	K (%)
T <sub>1</sub>	18.21	8.82	7.95	2.98	1.72	1.33
T <sub>2</sub>	17.23	8.09	7.11	2.86	1.60	1.31
T <sub>3</sub>	16.77	6.69	6.36	2.68	1.40	1.30
T <sub>4</sub>	17.71	8.47	7.72	2.90	1.68	1.33
T <sub>5</sub>	17.12	7.98	7.14	2.74	1.52	1.30
T <sub>6</sub>	16.61	6.99	6.85	2.66	1.47	1.30
T <sub>7</sub>	16.25	5.79	5.60	2.67	1.35	1.22
T <sub>8</sub>	15.61	5.23	5.32	2.62	0.95	0.91
F-test	*	*	*	*	*	*
SEm±	0.150	0.464	0.155	0.038	0.069	0.025
LSD <sub>5%</sub>	0.456	1.408	0.471	0.114	0.209	0.075
CV (%)	1.536	11.076	3.978	2.364	8.178	3.426

\*Significant at 5 % level of significance; \*\*T<sub>1</sub>. Standard check (RNPK + RFYM), T<sub>2</sub>. 75% NP through IF and 25% through *A. awamori* and *Azotobacter* sp. + RFYM + RK T<sub>3</sub>– 50% NP through IF and 50% NP through *A. awamori* and *Azotobacter* sp. + RFYM + RK, T<sub>4</sub> - T<sub>2</sub> + *T. harzianum*, T<sub>5</sub> – T<sub>3</sub> + *T. harzianum*, T<sub>6</sub> – RNPK from FYM, T<sub>7</sub> – RNPK only from IF, T<sub>8</sub> – control.

biofertilizers were used with reduced dosages of chemical fertilizers. Further, Mamatha and Bagyaraj (1999) stated the use of vesicular-arbuscular mycorrhizal (VAM) fungus, *G. fasciculatum* and mycorrhiza helper bacteria, *B. coagulans* in mulberry can reduce the use of IF by 50% without affecting both the leaf yield and quality variables.

### Nitrogen-phosphorus-potassium (NPK) percentage of mulberry leaf

Statistically significant ( $P < 0.05$ ) differences with respect to the NPK content of M<sub>5</sub> mulberry leaf was recorded among the different treatments (Table 4). Maximum leaf NPK of 2.98, 1.72 and 1.33%, respectively were recorded from T<sub>1</sub>, which was on par with T<sub>4</sub> (2.90, 1.68 and 1.33%). On the contrary, T<sub>8</sub> had the lowest leaf NPK of 2.62, 0.95 and 0.91%. The improvement in the leaf NPK percentage in mulberry leaves may be due to better fixation of 'N' by *Azotobacter*; mobilization and solubilization of P by *A. awamori* and *T. harzianum* in turn leading to better availability and absorption of the nutrients to plant system and its translocation to leaves. The improvement in K content of the leaf may be due to adequate availability of potassium through FYM and IF, particularly the recommended K in all the treatments except the control, and the synergistic effect of the inputs utilized.

The present results are in accordance with the findings of Das et al. (1994), who reported higher total nitrogen content in mulberry leaf when mulberry plant is inoculated with *Azotobacter*. Further, Krishnababu (1994) reported that there is an increase in NPK content of mulberry plant

applied with integrated chemical and bio-fertilizers. Moreover, Sukumar et al. (2001) concluded better efficiency in N and P uptake by mulberry when inoculated with *B. megaterium* and *A. chroococum* along with the recommended dosage of fertilizers. Increased uptake of NPK by mulberry as a result of integration of biofertilizers with different organic manures and inorganic fertilizers has been reported by Murali et al. (2006).

### Conclusion

In conclusion, 25% application of inorganic fertilizers (NP) to M<sub>5</sub> mulberry under rain-fed conditions can be reduced without affecting mulberry fresh leaf yield and quality variables by supplementing mulberry garden with integrated use of inorganic fertilizers, farm yard manure and bio-inoculants.

### ACKNOWLEDGMENTS

The authors express their sincere gratitude to Ministry of Education, Ethiopian Government's for the financial support without which this study would not been materialized. The technical support obtained from the staff of Department of Sericulture, University of Agricultural sciences, GKVK (Bangalore) is also highly appreciated.

### REFERENCES

A.O.A.C. (1970). Official Methods of Analysis. A.O.A.C., Washington, D.C., p. 105.

- Arnon DI (1949). Copper enzymes in isolated chloroplasts - Polyphenoloxidases in *Beta vulgaris*. Plant Physiol. 24:1-15.
- Bongale UD (2006). Soil Health Management for Sustainable Sericulture. National Seminar on Soil Health and Water Management for Sustainable Sericulture, Regional Sericulture Research Station (CSB), Bangalore, India 27-28 Sep. pp. 36-40.
- Cochran WG, Cox GM (2000). Experimental Design procedures for the behavioural Sciences. Cole publishing company, pp. 319-380.
- Dandin SB, Jayaswal J, Giridhar K (2003). Handbook of Sericulture Technologies, CSB, Bangalore, p. 259.
- Das PK, Choudhary PC, Ghosh A, Katiyar RS, Madhava Rao YR, Mathur VB, Mazumder MK (1994). Studies on the effect of bacterial bio-fertilizers in irrigated mulberry. Indian J. Seric. 33:170-173.
- Dubios MK, Gills A, Hamilton JK, Robeos RA, Smith R (1956). A calorimetric determination of sugars and related substances. Annal. Chem. 28:350-356.
- Hiscox JD, Israelstam GF (1979). A method for the extraction of chlorophyll from leaf tissue without maceration. Can. J. Bot. 57:1332-1334.
- Katiyar RS, Das PK, Choudhury PC, Ghosh A, Singh GB, Datta RK (1995). Response of irrigated mulberry (*Morus alba* L.) to VA-mycorrhizal inoculation under graded dosages of phosphorus. Plant. Soil 170:331-337.
- Krishnababu KS (1994). Influence of sources of phosphorus and P solubilizing fungus (*Aspergillus awamori*) on mulberry and cocoon production. M.Sc. (Seri.) Thesis, Univ. Agric. Sci., Bangalore, p.150.
- Lowry OH, Rosebrough N, Farr A, Randall R (1951). Protein measurement with Folin phenol reagent. J. Biol. Chem. 193:265-275.
- Mamatha RK, Bagyaraj DJ (1999). Response of already established mulberry in the field to inoculation with VA Mycorrhizal fungi and mycorrhiza helper bacteria. National Seminar on Tropical Sericulture University of Agricultural Sciences, Bangalore, India, p. 48.
- Murali C, Sreeramulu KR, Narayanaswamy TK, Shankar MA, Sreekantaiah M (2006). Effect of bio-inoculants and organic manures on soil microflora and fertility status of S<sub>36</sub> mulberry garden, National Seminar on Soil Health and Water Management for Sustainable Sericulture, Regional Sericultural Research Station, p. 90.
- Patil VC, Angadi SA, Roodagi LI, Angadi SS, Patil RR (2001). Response of mulberry to fertigation of N, P levels and mycorrhizal inoculation. Proc. Natl. Sem. Mulb. Seric. Res. KSSR & DI, Bangalore-India, pp. 375-378.
- Shankar MA, Devaiah MC, Anitha P, Rangaswamy BT (2000). Effect of graded levels of organic manures on growth, yield and quality of mulberry in relation to silkworm growth and cocoon production. Crop Res. 19:128-132.
- Shivakumar HR, Nageshchandra BK, Nagarajaiah C, Jagadish KS (2000). Impact of combined use of organic manures and inorganic fertilizers on growth, leaf yield and quality of mulberry. Proceedings of the National Seminar on Tropical Seric-Moriculture in Tropics, In: Chinnaswamy KP, Govindan R, Krishnaprasad NK, Reddy DNR (eds), UAS, Bangalore 1:94-96.
- Singhal BK, Chakraborti S, Mala VR, Sarkar A, Datta RK (2000). Photosynthesis for crop improvement in mulberry (*Morus* sp.). Sericologia 40:27-55.
- Singhal BK, Malav R, Sarkar A, Datta RK (1999). Nutritional disorders of mulberry (*Morus* spp.): III. Leaf nutrient guide for secondary nutrients. Sericologia 39(40):599-609.
- Singhvi NR, Kodandaramaiah J, Kumar JS, Singh GB (2007). Impact of feeding nutrient deficient mulberry leaves on cocoon production in silkworm, *Bombyx mori* L. J. Exp. Zool. 10(1):261-263.
- Sreeramulu KR, Bhaskar RN, Narayanaswamy TK (2004). Rhizosphere microflora of mulberry varieties as influenced by soil moisture condition. Organic Seric. & Seri-Byproduct Utilization, Natl. Sem. Prosp. pp. 94-98.
- Subbaswamy MR, Reddy MM, Sinha AK (1994). Thank silt: A cheap manure to mulberry, Indian silk 32(10):10.
- Sukumar J, Padma SD, Bongale UD, Veeresh M (2001). Response of mulberry to co-inoculation with *Azotobacter chroococcum* and *Bacillus megaterium*. Proc. Natl. Sem. Mulb. Seric. Res., KSSR & DI, Bangalore-India, pp.388-391.
- Sunil SH (2005). Influence of Organic Based Nutrient Management on Biochemical Parameters of Mulberry Leaf and Silkworm, *Bombyx mori* L. M.Sc. (Agil. Biochem.) Thesis, UAS, Bangalore p.110.
- Umesh MD (1999). Response of RFS<sub>135</sub> and M<sub>5</sub> varieties to Azotobacter inoculation in relation to growth, yield of mulberry and cocoon production under dry-land Alfisols. M.Sc. (Agil. Mic.) Thesis, UAS, Bangalore. p. 98.