

Infection prevalence of hydatidosis (*Echinococcus granulosus*, Batsch, 1786) in domestic animals in Ethiopia: A synthesis report of previous surveys

Abebe Fromsa^{1*} and Yilma Jobre²

¹College of Agriculture and Veterinary Medicine, Jimma University, P.O.Box 307, Jimma, Ethiopia, abebe.merga@ju.edu.et; afromsa@yahoo.com

²Food and Agriculture Organization of the United Nations, Cairo, Egypt, yilma.jobre@fao.org

* Corresponding author abebe.merga@ju.edu.et; afromsa@yahoo.com

Abstract

Hydatidosis/echinococcosis (*Echinococcus granulosus*, Batsch, 1786) is considered to be a serious problem for both public health and the livestock economy in Ethiopia. The present paper reviews and summarizes available data on the disease. Abattoir survey data collected over a period of 15 years (1985-1999) were analyzed to assess the infection prevalence of hydatidosis in domestic animals in three different agroecological zones of Ethiopia. Accordingly, 8036/22,863 (35.15%) cattle, 768/6518 (11.78%) sheep, 36/1753 (4.9%) goats, 70/417 (16.79%) camels and 0/150 (0%) pigs slaughtered in 21 different abattoirs located in various parts of the country were found harbouring hydatid cysts. A statistical discernible significant difference ($p < 0.001$) was observed in the overall hydatidosis infection prevalence between the different species of animals, and in infection prevalence of bovine hydatidosis in the three agroecological zones where the slaughtered cattle were believed to originate from. Similarly, a statistically significant difference ($p < 0.01$) in the prevalence of ovine hydatidosis was observed between mid-altitude and lowland agroecologies. Conversely, there was no significant difference ($p > 0.20$) in the infection prevalence of caprine hydatidosis between mid-altitude and lowlands. The present study reconfirms that hydatid disease is widespread and highly prevalent in ruminant livestock in Ethiopia and warrants institution of a nation-wide control measures. Considerations on experiences of other countries with respect to the economic benefits that hydatid disease control programs may bring to livestock producers and combining such efforts with other zoonosis control schemes in view of the 'one health' initiative is worthy for animal health planners and policy decision-makers. The paper also serves as a quick reference source on hydatidosis in the country and basis for future studies.

Keywords: Hydatidosis, infection prevalence, domestic animals, agroecology, Ethiopia

Introduction

Hydatid Disease is the name given to the condition caused by the zoonotic tapeworm *Echinococcus granulosus*. The tapeworm spends most of its adult life in the intestine of its definitive host, namely canids and in particular the dog. The tapeworm eggs become voided in the canids' faeces and as a result of ingesting the eggs, infection passes to the intermediate host, commonly herbivores while grazing. However, humans can become accidentally infected and hydatid cysts may develop throughout the body. Therefore, cystic echinococcosis (CE) or hydatidosis is a disease caused by the metacestode stage of *Echinococcus granulosus*. The disease is not apparent to farmers but is of considerable economic and public health importance (Torgerson and Budke, 2003; Ahmadi and Meshkehkar, 2011). In farm animals, it causes considerable economic loss due to condemnation of edible organs, decreased meat and milk production, reduced hide and fleece value and decrease in fecundity (Polydorou, 1981; Romazanov, 1983).

The incidence of human hydatid disease in any country is closely related to the prevalence of the disease in domestic animals and is highest where there is a large dog population and high sheep production (Khuroo, 2002). The most frequent strain associated with human cystic echinococcosis (hydatidosis) appears to be the common sheep strain (G1) (Arambulo III, 1997; Torgerson and Budke, 2003; Wani *et al.*, 2007). Recent molecular characterization of human and animal *Echinococcus granulosus* isolates demonstrated that the 'camel' strain (G6) is also equally important source of infection to humans (Shahnazi *et al.*, 2011; Magambo, 2006). The possible occurrence of the disease in humans in Ethiopia was indicated earlier by Graber, 1978. Besides, clinical and serologic tests conducted among the Dassanetch and Nyangatom pastoralist tribes of the south western part of the country revealed the prevalence of 4.8% palpable abdominal cysts, 15% hepatomegaly and 31.7% positive hydatid skin test (Fuller and Diane, 1981). A prevalence of 0.5% - 0.7% was also reported in Hamar pastoralist tribes of southwest Ethiopia (Macpherson *et al.*, 1989; Klungsøyr, 1993). During 1995 and 2005, 234 patients were operated for hydatid disease at Tikur Anbessa Hospital in Addis Ababa (Mesfin Minas *et al.*, 2007). In addition, a retrospective survey conducted between 2002 and 2006 revealed the registration of 24 hydatidosis cases out of the total of 36,402 patients, giving a mean annual incidence of 2.3 human hydatid cases per 100,000 people per year in north-western Ethiopia (Nigatu Kebede *et al.*, 2010).

Data on the prevalence of hydatidosis in animals will also provide a reliable indicator of the extent of environmental contamination with eggs from definitive carnivore hosts. Several studies have been conducted on livestock hydatidosis in different parts of the Ethiopia, particularly through the externship program of the Faculty of Veterinary Medicine (FVM), Addis Ababa University (AAU). However, most of these works were unpublished and not readily available to the wider national and international readership (Getaw *et al.*, 2010; Lemu Golassa *et al.*, 2011). As hydatidosis is considered a serious problem for public health and the livestock economy in Ethiopia, the present paper reviews and summarizes available data on the disease in the country. The present meta-analysis on livestock hydatidosis is based on FVM/AAU surveys (1985-1999), which allowed the depiction of long-term infection prevalence of the disease in domestic animals in different agroecologic zones, hence provide a clearer picture of the magnitude of the problem to animal health planners and decision makers. In addition, the paper serves as a quick reference source on hydatidosis in the country and basis for future studies.

Materials and Methods

Database management

The present study is based on surveys carried out on hydatidosis by graduating students of FVM/AAU between the periods 1985-1999 in 21 slaughterhouses located in different parts of the country, namely Addis Ababa, Arba Minch, Assela, Bahr Dar, Bale Robe, Debre Berhan, Dire Dawa, Dollo Oddo, Gondar, Harrar, Jijiga, Jimma, Kombolcha, Konso, Mekele, Melgue Wondo, Nazareth, Negele, Nekempte, Wolisso and Wollaita Soddo (Fig. 1). Individual survey data were carefully compiled, database organized by location (assumed origin) and species of animals (bovine, ovine, caprine, dromedary and swine). Infection prevalence of hydatidosis and levels of edible organ involvement were then determined.

Prevalence study

Prevalence was depicted by species of animals and agroecology of the study areas (Fig. 2). The latter are conventionally classified as highland (>2500 masl), mid-altitude (2500-1500 masl) and lowland (<1500 masl) zones. Debre Birhan and Bale Robe abattoirs are supplied from the highland whereas Arbaminch, Dire Dawa, Dollo Oddo, Konso, Negelle and Jijiaga abattoirs from Lowland and the remaining abattoirs from mid-altitude. It is generally assumed that animals raised in a radius of 50km are the ones feeding the nearby slaughterhouses.

Even if there are animals brought from outside of a given agroecological zone where the abattoir is located, it's again assumed that they constitute a very small proportion, hence do not alter the inferences drawn from the results.

Statistical Analysis

Chi-Square (χ^2) statistics at $p < 0.05$ was employed to depict the presence of significant difference between the prevalence of hydatidosis in various species and three agro-ecological zones (Fig. 2).

Results

Prevalence of hydatidosis

The surveys considered in this study reported that a total of 22863 cattle were slaughtered in the 18 abattoirs. Of these, 8036 (35.15%) were found harboring hydatid cysts. Besides, in 6397 (27.98%), 4211 (18.42%), 392 (1.71%) and 190 (0.83%) of the cases, hydatid cysts were located in lungs, liver, spleen, kidneys and heart, respectively. High prevalence (>61%) was observed at Bale Robe and Assela abattoirs. Additionally, the recorded prevalence in Assela was higher in the most recent report of 1997 (72.44%) than in 1990 (54.84%) and 1992 (50.29%). However, no increment was seen in the prevalence of the most recent years from Bahir Dar, Gondar, Kombolcha and Wallaita Sodo. The lowest (7.2%) prevalence reported was from Debre Berhan slaughterhouse. In most abattoirs, hydatidosis prevalence exceeded 30% (Table 1).

A total of 6518 sheep were slaughtered in 9 abattoirs, of which 768 (11.78%) were found infected with hydatid cysts. Furthermore, 531 (8.15%) had hydatid cysts in the lungs, 483 (7.41%) in the liver, 21 (0.3%) in the spleen, 8 (0.12%) in the kidneys and hearts. In most of the abattoirs, the infection prevalence reached 10%. In Assela, the recorded prevalence was higher in the most recent report of 1997 (53.5%, 130/243) than in 1990 (2.18%, 3/137). Similarly, in Kombolcha, the recorded prevalence was higher in 1996 (15.1%, 93/617) than in 1989 (4.4%, 39/888). The aggregated highest prevalence (35%) was recorded in Assela and the lowest prevalence (6.63%) in Nazareth abattoirs (Table 2).

Of the examined 1753 goats from 7 abattoirs, hydatidosis was confirmed only in 86 (4.9%) of the cases. The highest prevalence (9.3%) was recorded in Konso abattoir. There were no positive cases recorded from Assela and Kombolcha abattoirs. In terms of organ involvement, 37 (2.1%), 60 (3.42%), 3 (0.17%)

and 1(0.06%) of the cysts occurred in lungs, liver, spleen, heart and kidneys, respectively (Table 3).

Five of the surveyed abattoirs were handling camels and another two were handling swine. A total of 417 camels were examined and 70 (16.79%) were found harbouring hydatid cysts in their visceral organs. Organs infected include 59 (14.15%) lung and 13 (3.12%) liver (Table 4). None of the 150 pigs (110 from Addis Ababa and 40 from Kombolcha abattoirs) were infected with hydatid cysts (Table 4)

Variation in hydatidosis infection prevalence

The highest prevalence (35.15%) was recorded in cattle followed by dromedary (16.79%), ovine (11.87%), caprine (4.9%) and swine (0%). Statistical analysis revealed the presence of a significant difference ($p < 0.001$) in hydatidosis infection prevalence between the different species of examined animals (Table 5).

Further analysis showed a significant variation ($p < 0.001$) in infection prevalence of bovine hydatidosis along altitude zones. The highest prevalence (40.08%) of bovine hydatidosis occurred in the highlands followed by mid-altitude (38.09%) and lowland (18.95%). There was no data on the occurrence of ovine and caprine hydatidosis in the highlands. Ovine hydatidosis was significantly higher in the mid-altitude zone ($p < 0.01$) than in lowlands. Conversely, no significant difference ($p > 0.20$) was depicted in the prevalence of caprine hydatidosis along altitude zones. All camels slaughtered in the five studied abattoirs were originated from lowland areas (Fig. 3).

Table 1. Infection prevalence of bovine hydatidosis and edible organs condemned (1985-1999) in 18 slaughterhouses located in different parts of Ethiopia (Source: FVM/AAU)

Slaughterhouse location	No of animals examined	No. (%) positive cases	No. of organs with hydatid cysts (condemned as unfit for human consumption)				No of studies considered	Reference (source)
			Lungs	Liver	Spleen	Kidney		
Addis Ababa	825	121(14.67)	103	73	6	-	1	Gemeda, 1988.
Assela	380	133(35.00)	108	99	3	2	2	Alemayehu, 1990; Fekadu, 1997
Dire Dawa	1515	150(9.90)	107	79	-	-	1	Daniel, 1996
Dire Dawa, Harrar & Jijiga	565	53(9.38)	41	56	-	1	1	Woubet, 1987
Gondar	623	54(8.67)	30	39	-	-	1	Tamene, 1986
Jimma	227	28(10.12)	16	21	-	-	1	Abduljewad, 1988
Kombolcha	1505	132(8.77)	72	66	4	3	2	Asrat, 1996; Yilkal, 1989
Konso	345	65(18.80)	35	41	1	-	1	Fikre, 1994
Nazareth	483	32(6.63)	19	9	4	1	3	Yemane, 1990
Total No. (%)	6468	768 (11.87)	(8.15)	(7.41)	(0.3)	(0.12)	11	(0.12)

Table 2. Infection prevalence of ovine hydatidosis and edible organs condemned (1985-1999) in 9 surveyed abattoirs located in different parts of Ethiopia (Source: FVM/AAU)

Slaughterhouse location	No of animals examined	No. (%) positive cases	No. of organs with hydatid cysts (condemned as unfit for human consumption)					No of surveys considered	References
			Lungs	Liver	Spleen	Kidney	Heart		
Addis Ababa	1027	217(21.13)	201	94	4	18	6	1	Gemeda, 1988
Arbaminch	568	147(25.88)	124	64	1	4	1	1	Mohammed, 1988
Assela	2325	1430(61.51)	1223	1032	88	95	35	3	Alemayehu, 1990; Dessie, 1992; Fekadu, 1997
Awassa	1020	350(34.30)	270	81	19	2	-	1	Getachew, 1991
Bahr Dar	1907	851(44.63)	613	403	23	63	50	2	Musie, 1995; Nebiyou, 1990
Bale Robe	764	481(62.96)	302	240	45	25	21	1	Woubet, 1988
Debre Berhan	531	38(7.2 0)	23	17	1	4	2	1	Tsegaye, 1995
Dire Dawa	2224	305(13.70)	204	241	-	-	-	1	Dainel, 1996
Dire Dawa, Harrar and Jijiga	854	239(27.99)	170	134	13	6	4	1	Woubet, 1987
Gonder	1697	478(28.17)	366	203	18	6	12	2	Roman, 1987; Tamene, 1986
Jimma	1010	340(33.66)	268	79	16	1	-	1	Abduljewad, 1988
Kombolicha	1670	599(35.87)	467	337	33	9	21	2	Asrat, 1996; Yilkal, 1989
Konso	447	115(25.70)	52	93	1	1	-	1	Fikre, 1994.
Mekele	548	176(32.12)	144	95	8	-	1	1	Hagos, 1997.
Melgue wondo	1878	807(42.90)	670	481	53	11	9	1	Getahun, 1987
Nekemet	1978	617(31.19)	572	238	4	19	7	1	Feyessa, 1987
Nazareth	1185	447(37.72)	424	205	33	10	21	1	Yemane, 1990
Wallaita	1230	399(32.43)	304	174	32	6	3	2	Abdul, 1992, Abel, 1985
Total (%)	22863	8036 (35.15)	6397 (27.98)	4211 (18.42)	392 (1.71)	280 (1.22)	190 (0.83)	24	

Table 3. Prevalence of caprine hydatidosis and edible organs condemned (1985-1999) in 7 surveyed abattoirs located in different parts of Ethiopia (Source: FVM/AAU)

Slaughterhouse location	No of animals examined	No. (%) positive cases	No. of organs with hydatid cysts (condemned as unfit for human consumption)				No of studies considered	Reference (source)
			Lung	Liver	Spleen	Kidney		
Addis Abeba	170	12(7.05)	4	11	-	-	1	Gemeda, 1988.
Assela	124	0(0)	-	-	-	-	2	Alemayehu, 1990 & Fekadu, 1997
Dire Dawa, Harrar & Jijiga	184	12(6.50)	4	10	-	-	1	Woubet, 1987
Gonder	194	9(4.64)	5	7	-	-	1	Tamene, 1986
Kombolcha	60	0(0)	-	-	-	-	1	Asrat, 1996
Konso	389	36(9.30)	19	21	-	-	1	Fikre, 1994.
Nazareth	632	17(2.7)	5	11	3	2	1	Yemane, 1990
Total	1753	86(4.9)	37 (2.1%)	60 (3.42%)	3 (0.17%)	2 (3.11%)	8	

Table 4. Prevalence of hydatidosis in swine and camel and edible organs condemned (1985-1999) in 4 surveyed abattoirs located in different parts of Ethiopia (Source: FVM/AAU)

Slaughterhouse location	Species examined	No of animals examined	No. (%) positive cases	No. of organs with hydatid cysts (condemned as unfit for human consumption)				No of studies considered	Reference (source)
				Lungs	Liver	Spleen	Kidney		
Addis Ababa	Swine	110	0(0%)	-	-	-	-	1	Gemeda, 1988.
Kombolicha	Swine	40	0(0%)	-	-	-	-	1	Yilkal, 1989
Dire Dawa, Harrar & Jijiga	Dromedary	296	43(14.53%)	36	8	-	-	2	Ahemed, 1998; Woubet, 1987
Dollo Oddo and Nagelle	Dromedary	121	27(22.3%)	23	5	-	-	1	Balako, 1999
Total	Dromedary	417	70 (16.79%)	59 (14.15%)	13 (3.12%)	-	-	3	
	Swine	150	0(0%)	-	-	-	-	2	

Table 5. Prevalence of hydatidosis in different species of examined animals

Animal species	No. examined	No. (%) positive	No. of studies considered	χ^2 (p-value)
Bovine	22863	8036 (35.15)	24	1963 (<0.001)
Ovine	6468	768 (11.87)	11	
Caprine	1753	86 (4.90)	7	
Dromedary	417	70 (16.79)	3	
Swine	150	0 (0)	2	



Figure 1: Map of Ethiopia showing the locations of abattoirs where the surveys were carried out.




Figure 2. Elevation map of Ethiopia showing altitude ranges from -121m below sea level to over 4000m above sea level (Source: Berhanu Bedane)




Figure. 3. Infection prevalence of hydatidosis in different animal species versus altitude zones.

Discussion

Echinococcosis is one of the most geographically widespread zoonotic diseases that occur in all inhabited continents, including sub-arctic, arctic, temperate, subtropical and tropical zones, especially in undeveloped and developing countries. The disease is endemic to hyperendemic in agricultural countries of Europe, northern, eastern and southern Africa, southern and northern America, Middle East and Asia (Matossian, *et al.*, 1977; Wen and Yang, 1997; Arambulo III, 1997; Torgerson and Budke, 2003; Budke *et al.*, 2006; Dakkak, 2010). In Ethiopia, the Faculty of Veterinary Medicine of Addis Ababa University served as one of the leading organizations in collecting baseline information on various animal diseases in the country. As a result, it has accumulated a wealth of information on the status of hydatidosis in animals collected from various parts of the country. However, most of this information is unavailable to wider national and international readership. Consequently, in an attempt to fill-in the prevailing information gap, the present work reviewed and summarized the abattoir survey data, collected over a period of 15 years (1985-1999) through the externship programme of the Faculty of Veterinary Medicine, Addis Ababa University.

The analysis revealed a hydatidosis prevalence rate of 34.15% in cattle, 11.87% in sheep, 4.9% in goats, 16.79% in camels and 0% in pigs. In Assela, where there are multiple records, higher prevalence of bovine and ovine hydatidosis was observed in the most recent report than in the older ones. The mean prevalence value reported in the present work concurs with the recently reported prevalence of 34.05% in cattle and 10.6% in sheep from Bahir Dar (Nigatu Kebede *et al.*, 2009c), 32.11% in cattle from Mekelle (Gebretsadik Berhe *et al.*, 2010) and 29.69% in cattle from Ambo (Endrias Zewdu *et al.*, 2010). Slightly higher prevalence of 48.9% in cattle from Debre Markos (Kebede Nigatu, *et al.*, 2009a), 46.8% in cattle, 29.3% in sheep, and 6.7% in goats from Nazareth (Getaw *et al.*, 2010) and 52.69% in cattle from Hawassa (Feyessa Regassa *et al.*, 2010) were also reported. A retrospective survey of bovine hydatidosis conducted in Gondar, Injibara and Finote Selam municipal abattoirs during 2002 to 2007 revealed an increasing trend from year to year in prevalence of bovine hydatidosis that caused the condemnation of 79.5% organs (Nigatu Kebede, 2010). This might be attributed to backyard slaughter practice, an increase in the population of stray dogs and the absence of control program. Bovine hydatidosis prevalence of 22.1% was reported from Tigray region (Kebede, *et al.*, 2009), 13.86% from 7 export abattoirs (Solomon Hailemariam, 1975), 15.2% from Birre-Sheleko and Dangila abattoirs (Nigatu Kebede *et al.*, 2011), and 16.85%, 16.0% and 15.4% from Wolayita Sodo by Jemere Bekele and Berhanu Butako (2011), Nigatu Kebede *et al.*, (2009b) and Regassa, *et al.*, (2009) respectively. Varying prevalence rates were also reported by different authors from diverse countries in different intermediate hosts. For example, an infection prevalence of 48% in cattle, 63.8% in sheep and 34.7% in goat from Ngorongoro district (Ernest *et al.*, 2004) and 6.02% in shoats and 4.2% in cattle from Arusha area of Tanzania (Nonga and Karimuribo, 2009), 19.4% in cattle, 3.6% in sheep, 4.5% in goats and 61.4% in camels from Kenya (Njoroge *et al.*, 2002), 6.6% in cattle, 4.3% in sheep and 27.2% in camels from Libya (Mohammed, 1985), 31.6% in cattle, 24.4% in sheep, 42.2% in goats and 59.9% in pigs from Niger (Arene, 1985), 8.28% in cattle, 12.61% in sheep and 6.56% in goats and 32.85% in camels from Saudi Arabia (Ibrahim, 2010), 3% in cattle, 7% in sheep and 45% in camels from Sudan (Elmahdi *et al.*, 2004) and 18.3% in sheep from Argentina (Larrieu *et al.*, 2001). Sotiraki *et al.*, (2003) reported a prevalence of 82% in cattle, 80% in sheep, 24% in goats and 5% in pigs from Greece before the introduction of control program in 1984 which reduced the occurrence of the disease to 0% in cattle, to 31.3% in sheep, to 10.3% in goats and to 0.6% in pigs. A decrease in annual prevalence of camel liver hydatidosis from 24.1% in 2004 to 6.8% in 2010 and a corresponding decrease in annual

prevalence of camel lung hydatidosis from 28.7% in 2004 to 7.1% in 2010 were also reported from Iran as a result of greater awareness created about echinococcosis among farmers (Borji, 2011).

The difference in infection prevalence rate between countries could be associated with different factors like control measures put in place, the level of community awareness created about the disease, education and economic status of the population and the farming community. The absence of hydatid cyst infection in pigs in Ethiopia could be mainly attributed to the absence of extensive swine farms in the country. All farms are under intensive husbandry practices where farm facilities are fenced and animals provided with concentrate feed and industrial byproducts.

The result of this study revealed the existence of statistically significant association ($p < 0.001$) between bovine/ovine hydatidosis prevalence rates and the altitude zones, the highest prevalence reported in highlands followed by mid altitude zones. This difference might be ascribed to the differences in the socio-economic factors and animal husbandry practices that may facilitate close contacts between domestic animals and village dogs, the latter being allowed free access to hydatid infected offals. The distribution may also be influenced by temperature and humidity (Wen and Yang, 1997). Similar spatial variation in the infection prevalence in livestock was previously reported by Yilma Jobre *et al.*, (1996); Kebede, *et al.*, (2009); and Ibrahim, (2010). Caprine hydatidosis prevalence rates of 4.72% and 6.5% were recorded in mid-altitude and lowland areas, respectively. There was no statistically significant difference ($p > 0.20$) in hydatidosis prevalence between the two agroecological zones.

The distribution may also be influenced by

Statistical analysis of hydatidosis infection rates on the basis of species, indicated that there exists a significant difference ($p < 0.001$) between cattle, sheep, goats, camels and pigs. This finding is in agreement with previous study by Yilma Jobre *et al.*, (1996) and Nigatu Kebede *et al.*, (2009c). The difference in infection rates between different species of animals might be explained mainly as a result of the involvement of different strains of *E. granulosus*, variations in feeding behaviors of the animals and animal husbandry practices. There is an evidence that indicates the existence of a number of strains of *E. granulosus*, which differ morphologically and biochemically. These include strains, which utilize sheep/dog, horse/dog, Camel/dog, pig/dog, buffalo/dog, goat/dog, cattle/dog and man/dog cycles (Soulsby, 1986). High prevalence rate of hydatidosis

in cattle and camel in Ethiopia could be ascribed to age factor and method of husbandry. Cattle and camels are generally slaughtered at an older age than sheep and goats and consequently are exposed to infection over a longer period of time. Ibrahim (2010) and Cabrera *et al.*, (2003) reported an increase in prevalence of the disease with increase in age. Study conducted in Uruguay by Guorino *et al.*, (1981) also showed the impact of age on the incidence of the disease and an increase in prevalence of the disease (reaching 46-50%) in older subjects as compared to 9.4% in lambs. In the same study a prevalence rate of 81.2% in bulls and older bullocks, 69.7% in fattening bullocks and 51.1% in cows were reported in contrast to 7% in calve. A survey made in New Zealand also showed that the incidence increased with age from 9% in cattle of 1-2 years old to 74% in cattle of 5 years or older (Gracy, 1994).

This report indicated that lungs and liver were the most commonly affected organs by hydatid cysts in cattle, sheep, goats and camels. This is in fundamental agreement with literature, which states that hydatid cysts are most commonly found in the liver and lungs of ungulates (Soulsby, 1986; Urguhart *et al.*, 1988; Yilma Jobre *et al.*, 1996; Getaw *et al.*, 2010; Ibrahim, 2010; Nigatu Kebede, 2010). This could be justified by the fact that lungs and liver possess the first greater capillary fields which acts as partial barriers for the ingested oncosphere which adopt the portal vein route and primarily negotiates the hepatic and the pulmonary filtering action sequentially before any other peripheral organ is invaded (Matosain, 1977). In this report percent involvement of the lungs is higher than the liver in cattle, sheep and camel and the vice versa holds true in goats. Literature reveals that in older cattle and sheep, the liver sinusoids permits the embryos to traverse the sinusoids and reach the lungs. It is also possible for embryos to enter the hepatic circulation and be carried, via the thoracic duct and heart, to the lungs so that lungs will be infected before or instead of the liver (Gracy, 1994). The same may hold true in case of camels. In goats the sinusoids of the liver might be narrow so that the majority of hydatid embryos are arrested in that organ. Similar higher percent involvement of the liver than the lungs was recorded in pigs and humans (Wen and Yang, 1997; Khuroo 2002; Torgerson and Budke, 2003).

There is no doubt that the Faculty of Veterinary Medicine of the Addis Ababa University has made extensive studies and serves as repository of baseline information for the country. However, there are obvious gaps in terms of geographic coverage. For instance, no studies on the prevalence of hydatidosis were carried out in Afar, Beneshangul Gumuz and Gambela. The different veterinary schools should consider the case of hydatidosis as just one example

and plan to undertake a harmonized and well plan animal health investigation plans in such a way that the veterinary authorities will make good use of the research outputs in animal health programs planning and decision-making.

The control of diseases such as hydatidosis must consider its economic feasibility of intervention without forgetting the huge direct and indirect loss due to it. In addition, possibility of amalgamating hydatid disease control planning with other zoonotic disease control programs is essential. The current FAO initiative – one health – approach for sustainable animal health contained animal-related human health risks is worth while considering. FAO’s strategic vision is “a world in which risks to animal and animal-related human health due to a wide range of high-impact emerging and re-emerging zoonotic and non-zoonotic diseases, and their associated impacts on food security, livelihoods, trade and economic development are minimized through prevention, early detection, rapid response, containment and elimination. The main goal of the strategy is to establish a robust global animal health system that effectively manages major animal health risks, paying particular attention to the animal-human-ecosystem interface, and placing disease dynamics into the broader context of agriculture and socio-economic development and environmental sustainability” (FAO, 2011).

References

- Abdul, J.R, 1992. Economic significance of bovine fasciolosis and hydatidosis at Soddo abattor, DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp 23.
- Abduljewad, A, 1988. Hydatidosis prevalence at Jimma abattoir. DVM Thesis. Addis Ababa University, Faculty Of Veterinary Medicine, Debre Zeit, Ethiopia, Pp25.
- Abel, M, 1985. A survey of econococcus and its public health significance in Soddo abattoir, DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp28.
- Ahemed, I.D., 1998. A preliminary investigation on major disease of camels in Eastern Ethiopia: abattoir and field survey. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp45.
- Ahmadi, N.A. and Meshkehkar, M., 2011. An abattoir-based study on the prevalence and economic losses due to cystic echinococcosis in slaughtered herbivores in Ahwaz, south-western Iran. *J. Helminthol.* 85, 33–39.

- Alemayehu, L., 1990. The prevalence of hydatidosis in cattle, sheep and goats and *echinococcus granulosus* in dogs in Arsi administrative region, DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp26.
- Arambulo III, P., 1997. Public health importance of cystic echinococcosis in Latin America. *Acta Trop.*, 67, 113-124.
- Arene, F.O.I., 1985. Prevalence of hydatid cysts in domestic livestock in the Niger Delta. *Trop. Anim. Hlth. Prod.*, 17, 3-5.
- Asrat, G., 1996. Prevalence and economic significance of hydatidosis or echinococcosis in slaughtered cattle, sheep and goats in South Wollo. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp36.
- Balako. G., 1999. Observation on disease of one- humped Camel in southern Ethiopia: abattoir survey. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp34.
- Bekele, J. and Butako, B., 2011. Occurrence and financial loss assessment of cystic echinococcosis (hydatidosis) in cattle slaughtered at Wolayita Sodo municipal abattoir, Southern Ethiopia . *Trop. Anim. Hlth. and Prod.*, 43(1), 221-228
- Berhe, G., Tadesse, G, Kiros, H. and Negus, A., 2010. Concurrent infection of hydatidosis and fasciolosis in cattle slaughtered at Mekelle Municipal abattoir, Tigray Region. *Ethiop. Vet. J.* 14(2), 39-49.
- Bersissa, K., 1994. Hydatidosis in Nekempte: prevalence in slaughtered cattle and sheep, estimated economic loss and incidence in stray dogs. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp15.
- Borji, H., Azizzadeh, M. and Afsai A., 2011. An abattoir-based study of hydatidosis in the dromedary (*Camelus dromedarius*) in Mashhad, Iran. *J. Helminthol.*, 1-2.
- Budke, C.M., Deplazes, P. and Torgerson P.R., 2006. Global socioeconomic impact of cystic echinococcosis. *Emerging Infectious Diseases.* 12, 296-303.
- Cabrera, P.A., Irabedra, P., Orlando, D., Rista, L., Hara'n, G., Vin'als, G. and Blanco, M.T. et. al., 2003. National prevalence of larval echinococcosis in sheep in slaughtering plants *Ovis aries* as an indicator in control programmes in Uruguay. *Acta Trop.*, 85, 281-285.
- Dainel, F., 1996. Economic importance of organ condemnation due to fasciolosis and hydatidosis in cattle and sheep slaughtered at Dire Dawa abattoir. DVM Thesis.

- Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp30.
- Dakkak, A., 2010. Echinococcosis/hydatidosis: A severe threat in Mediterranean countries. *Vet. Parasitol.*, 174, 2–11
- Dessie, S., 1992. Economic significant of bovine hydatidosis, fasciolosis and cysticercosis. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp30.
- Dinka, A., 1996. Prevalance and economic significance of hydatidosis or echinocooccosis in slaughtered cattle, sheep and goats in south Wollo. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp39.
- Elmahdi, I.E., Ali, Q.M., Magzoub, M.M., Ali Abraham, A.M., Saad, M.B., Romig, T., 2004. Cystic echinococcosis of livestock and humans in central Sudan. *Ann. Trop. Med. Parasitol.*, 98, 473–479.
- Ernest, E. Kassuku, A., Kazwala, R., 2004. Studies on the epidemiology of echinococcosis/ hydatidosis in Ngorongoro district, Arusha region, *Tanzania. Int. Arch. Hydatid.* 35,43.
- FAO, 2011. Sustainable animal health and contained animal-related human health risk – in support of the emerging ‘One Health’ agenda. Applying Lessons Learned from Highly Pathogenic Avian Influenza in the Prevention and Containment of Major Animal Diseases and Related Human Health Risks. Working document, March 21-25, 2011, Rome, Italy.
- Fekadu, O.A., 1997. Study on prevalence and economic significance of hydatidosis in ruminants and *echinococcus granulosus* in dogs. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp49.
- Feyessa, R., 1987. Prevalence of hydatidosis in Nekempte municipality slaughterhouse. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit Ethiopia, pp26.
- Fikre, L., 1994. Echinococcosis/hydatidosis in Konso (southern Ethiopia) an assessment trail of its prevalence, economic and public health importance. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp42.
- Fuller, G.K. and Diane, C.F., 1981. Hydatid disease in Ethiopia clinical survey with some immunodiagnostic test results. *Am. J. Trop. Med. Hyg.*, 30(3), 645-652.

- Gemeda, B., 1988. A preliminary survey on echinococcosis/hydatidosis in livestock (cattle, sheep, goats, and pig) slaughtered at Addis Ababa abattoir. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp23.
- Getachew, J., 1991. The prevalence of hydatidosis in cattle at Awassa abattoir. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp23.
- Getahun, D., 1987. Incidence of bovine echinococcosis at Melgue Wondo abattoir and the role of dogs, jackals and hyena in the transmission around Awassa and Wondogenet. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp21.
- Getaw, A., Beyene, D., Ayana D., Megersa B., Abunna, F., 2010. Hydatidosis: Prevalence and its economic importance in ruminants slaughtered at Adama municipal abattoir, Central Oromia, Ethiopia. *Acta Tropica.*, 113, 221–225.
- Golassa, L., Abebe, T. and Hailu, A., 2011. Evaluation of crude hydatid cyst fluid antigens for the serological diagnosis of hydatidosis in cattle. *J. Helminthol.* 85, 100–108.
- Graber, M., 1978. Helminth and Helminthiasis of domestic and wild animals in Ethiopia. *Revue Elev. Med. Vet. Pays Trop.*, 1, 13-95.
- Gracy, J.F., 1994. Textbook of meat hygiene. 6th ed., Beilleir Tindall, London, pp517.
- Guorino, M.L.A., Cardoso De Ghi, M. Clivio De Freire, S., Freire, H., Gaudiano, M.J., and La Gamma. G., 1981. Control of hydatidosis in Uruguay. *Vet. Med. Rev.*, 1, 47-57.
- Hagos, Y., 1997. Hydatidosis/echinococcosis: prevalence and economic impact in bovine at Mekele Municipal abattoir, zoonosis and infection in dogs (Mekele Tigray). DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp35.
- Ibrahim, M.M., 2010. Study of cystic echinococcosis in slaughtered animals in Al Baha region, Saudi Arabia: Interaction between some biotic and abiotic factors. *Acta Trop.*, 113, 26–33.

- Jobre Y, Lobago F, Tiruneh R, Abebe G, Dorchies P., 1996. Hydatidosis in three selected regions in Ethiopia: an assessment trial on its prevalence, economic and public health importance. *Revue Méd. Vét.*, 147 : 797-804
- Kebede N, Mitiku A and Tilahun G., 2010. Retrospective survey of human hydatidosis in Bahir Dar, north-western Ethiopia. *East Mediterr Hlth J.*, 16, 937-41.
- Kebede, N., 2010. A retrospective survey of bovine hydatidosis in three abattoirs of Amhara National Regional State, northwestern Ethiopia. *Trop. Anim. Health. Prod.*, 42,323–325
- Kebede, N., Abuhay, A. Tilahun, G. and Wossene, A., 2009a. Financial loss estimation, prevalence and characterization of hydatidosis of cattle slaughtered at Debre Markos Municipality abattoir, Ethiopia. *Trop. Anim. Health. Prod.* 41, 1787–1789.
- Kebede, N., Gebre-Egziabher, Z., Tilahun, G. and Wossene, A., 2011. Prevalence and Financial Effects of Hydatidosis in Cattle Slaughtered in Birre-Sheleko and Dangila Abattoirs, Northwestern Ethiopia. *Zoonoses Public Hlth.*, 58, 41–46.
- Kebede, N., Mekonnen, H., Wossene A. and Tilahun, G., 2009b. Hydatidosis of slaughtered cattle in Wolaita Sodo Abattoir, southern Ethiopia. *Trop. Anim. Hlth. and Prod.*, 41(4), 629-633
- Kebede, N., Mitiku, A. and Tilahun, G., 2009c. Hydatidosis of slaughtered animals in Bahir Dar Abattoir, Northwestern Ethiopia. *Trop. Anim. Health. Prod.*, 41, 43–50
- Kebede, W., Hagos, A., Girma Z. and Lobago, F., 2009. Echinococcosis/hydatidosis: its prevalence, economic and public health significance in Tigray region, North Ethiopia. *Trop. Anim. Hlth. and Prod.*, 41(6), 865-871.
- Khuroo, M.S., 2002. Hydatid disease: current status and recent advances. *Ann. Saudi Med.* 122, 56-64.
- Klungsoyr, P Courtright, P. and Hendrikson, T.H. (1993): Hydatid disease in the Hamar of Ethiopia: a public health problem for women. *Transact. Royal Soc. Trop. Med. Hyg.*, 87(3), 254-255.
- Larrieu, M., Costa, M.T, Cantoni, G., Alvarez , R., Cavagion, L., Labanchi, J.L., et al., 2001. Ovine Echinococcus granulosus transmission dynamics in the province of Rio Negro, Argentina, 1980–1999. *Vet. Parasitol.*, 98, 263–272.

- Macpherson, CNL., Spoerry, A., E. Zeyhle, E., Romig, T., Gorfe, M. (1989): Pastoralists and hydatid disease: an ultrasound scanning prevalence survey in East Africa. *Transact. Royal Soc. Trop. Med. Hyg.*, 83(2), 243-247.
- Magambo, J., Njoroge, E., Zeyhle, E., 2006. Epidemiology and control of echinococcosis in sub-Saharan Africa. *Parasitol. Int.* 55, 193-195.
- Matosain, R.M., 1977. Hydatidosis, A global problem of increasing importance. *Bul. WHO.* 55, 499-507.
- Minas, M., Biluts, H., Bekele, A. and Alemie, M., 2007. Surgical management of 234 patients with hydatid disease: the Tikur Anbessa Hospital experience. *Ethiop Med J.*, 45, 257-65.
- Mohammed, A, 1988. Study on prevalence and economic significance of bovine hydatidosis in Gamogofa region. DVM thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp29.
- Mohammed, A.A., 1985. Prevalence of *Echinococcus granulosus* among domestic animals in Libya. *Trop. Anim. Hlth. Prod.*, 17, 169-170.
- Musie, H., 1995. Bovine hydatid disease: an assessment trial of its prevalence and economic importance at Bahr Dar Municipality abattoir. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, Pp32.
- Nebiyou, G., 1990. Study of hydatidosis/echinococcosis in cattle slaughtered at Bahr Dar Municipality abattoir. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp24.
- Njoroge, E.M., Mbithi, P.M., Gathuma, J.M., Wachira, T.M., Magambo, J.K., Zeyhle, E.A., 2002. Study of cystic echinococcosis in slaughter animals in three selected areas of northern Turkana, Kenya. *Vet. Parasitol.*, 104, 85-91.
- Nonga, H. E. and Karimuribo E. D., 2009. A retrospective survey of hydatidosis in livestock in Arusha, Tanzania, based on abattoir data during 2005 – 2007. *Trop. Anim. Health. Prod.*, 41, 1253–1257
- Polydorou, K., 1981. Animal health and economics. Case study: echinococcosis with reference to Cyprus. *Bul. Int. Epz.*, 93, 981-992.
- Regassa, A., Abunna, F., Mulugeta A. and Megersa B., 2009. Major metacestodes in cattle slaughtered at Wolaita Soddo Municipal abattoir, southern Ethiopia:

- Prevalence, cyst viability, organ distribution and socioeconomic implications. *Trop. Anim. Hlth. and Prod.*, 41(7), 1495-1502
- Roman, T., 1987. Study on economic significance of bovine fascioosis and hydatidosis at Gondar abattoirs. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp36.
- Romazanov, V.T., 1983. Evaluation of economic losses due to echinococcosis. In: Lysendo A, editor. Zoonosis control: collection of teaching aids for international training course vol. II. Moscow: Centre of International Projects GKNT, pp 283–85.
- Shahnazi, M., Hejazi, H., Salehi, M. and Reza Andalib, A.A., 2011. Molecular characterization of human and animal *Echinococcus granulosus* isolates in Isfahan, Iran. *Acta Trop.*, 117, 47–50.
- Solomon, H.M., 1975 A brief analysis of activity of the meat inspection and quarantine division. Department of veterinary services, pp128.
- Sotiraki, S., Himonas, C. and Korkoliakou, P., 2003. Hydatidosis-echinococcosis in Greece. *Acta Trop.*, 85, 197-201
- Soulsby, E.J.L., 1986. Helminth, Arthropod and protozoa of domestic animals. 7th edition, Bailleir Tindall, London, UK, pp809.
- Tamene, M., 1986. A preliminary study on echinococcosis/hydatidosis of livestock (cattle, sheep and dogs) in Gondar administrative region. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp286.
- Torgerson, P.R. and Budke, C.M., 2003. Echinococcosis – an international public health challenge. *Res. Vet. Sci.*, 74, 191–202
- Tsegaye, T., 1995. Epidemiology of bovine fasciolosis and hydatidosis in Debre Berhan region. DVM Thesis, F.V.M., Debre Zeit, Ethiopia, pp56.
- Urguhart, G.M., Armour, J., Duncan, J.L., Dunn, A.M. and Jennings, F.W., 1988. Veterinary parasitology, Longman, UK, pp228.
- Wani, M.M., Durrani, A.M., Shafi, M., Wani Mubbashir, M. and Khan, M., 2007. Hydatid Disease of the Soft Tissues of the Lower Limb: case report. *JK-Practitioner.*, 14, 104-106.
- Wen, H. and Yang, W.G., 1997. Public health importance of cystic echinococcosis in China. *Acta Trop.*, 67, 133-145.

- Woubet, S., 1988. Prevalence of cattle hydatidosis and its economic significance in Robe municipality abattoir. DVM Thesis F.V.M., A.A.U., Debre Zeit, Ethiopia, pp35.
- Woubet. M., 1987. A preliminary study on echinococcosis/hydatidosis in Harrargie region and the efficacy of *Glinus Lotoides* seeds against *Echinococcus granulosus*, in pups infected experimentally with hydatid materials. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp30.
- Yemane, G., 1990. Preliminary study on echinococcosis/hydatidosis in ruminants slaughtered at Nazareth abattoir, D.V.M Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp27.
- Yilkal, A., 1989. Hydatidosis in cattle, sheep, pigs, *Cysticercus tenuicollis* in sheep around Dessie and efficacy of *Hagenia abyssinica* (Kosso) on *Taenia hydatigena*. DVM Thesis. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp27.
- Zewdu, E., Teshome, Y. and Wakwoya, A., 2010. Bovine hydatidosis in Ambo Municipality abattoir, West Shoa, Ethiopia. *Ethiop. Vet. J.*, 14, 1-14.

