



JIMMA UNIVERSITY

SPORT ACADEMY

DEPARTMENT OF SPORT SCIENCE

**EFFECTS OF 12 WEEKS OF AEROBIC EXERCISE PROGRAM ON WEIGHT LOSS
OF SEDENTARY OVERWEIGHT WOMEN IN SHESHEMANNE CITY**

BY: UKULA YOHANNIS

**A RESEARCH SUBMITTED TO THE SCHOOL OF GRADUATE STUDIES OF JIMMA
UNIVERSITY SPORT ACADEMY DEPARTMENT OF SPORT SCIENCE IN PARTIAL
FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF
EDUCATION (MED).**

SEPTEMBER, 2021

JIMMA, ETHIOPIA

JIMMA UNIVERSITY
SPORT ACADEMY
DEPARTMENT OF SPORT SCIENCE

**EFFECTS OF 12 WEEKS OF AEROBIC EXERCISE PROGRAM ON WEIGHT LOSS
OF SEDENTARY OVERWEIGHT WOMEN IN SHESHEMANNE CITY**

BY: -UKULA YOHANNIS

ADVISO: WONDEMAGEGN DEMISE (PhD)

CO-ADVISO: SZEWGE HAILU (MSc)

SEPTEMBER, 2021

JIMMA, ETHIOPIA

DECLARATION

JIMMA UNIVERSITY

SPORT ACADEMY

DEPARTMENT OF SPORT SCIENCE

I, Ukula Yohannis, hereby declare that “The Effects of 12 weeks of aerobic exercises program on weight loss of sedentary overweight women in Shashemene City” is my original work which has not been presented for any degree or examination in any University and the sources used have been duly acknowledged and cited.

This thesis has been submitted in partial fulfillment of the requirements for Master of Science. Degree in Sport Science at Jemma University and is deposited at the University Library to be made available to borrowers under rules of the Library. I solemnly declare that this thesis is not submitted to any other institution anywhere for the award of any Academic Degree, Diploma, or Certificate. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgement of source is made. Requests for permission for extended quotation from or reproduction of this manuscript in whole or in part may be granted by the Head of the Major Department or Graduate Studies when in his or her judgment the proposed use of the material is in the interests of scholarship. In all other instances, however, permission must be obtained from the Author.

By: Ukula Yohannis

Sign _____ Date _____

Advisor: Wondimegegh Demise

Sign _____ Date _____

Co-Advisor: Zewuge Hailu

Sign _____ Date _____

APPROVAL SHEET

JIMMAUNIVERSITY

SPORT ACADEMY

DEPARTMENT OF SPORT SCIENCE

As a thesis advisor, I hereby certify that I have read and evaluate this thesis entitled the effects of 12 weeks aerobic exercise program on weight loss of sedentary overweight women in Sheshemenne city prepared under my guidance by Ukula Yohannis I recommend that to be submitted as fulfilling the thesis requirement.

Submitted by: Ukula Yohannis

Signature _____ Date _____

Name of Advisor: Wondimegegh Demise

Signature _____ Date _____

Name of Co-Advisor: Zewuge Hailu

Signature _____ Date _____

Name of internal examiner: _____

Signature _____ Date _____

Name of external examiner: _____

Signature _____ Date _____

Graduate Thesis Ownership Agreement

This thesis is a property of Jimma University, an institution that awarded Degree to the graduate student and funded its research cost fully or partly. The research work was accomplished under the close support and supervision of the assigned University's academic staff. It is therefore strictly forbidden to publish, modify, or communicate to or put at the disposal of third party the entire document or any part therefore without the common consent of the research supervisor(s) and the graduate student. Disregarding this agreement would lead to accountability according to the Jimma University's Research and Publication Misconduct Policy Article 1.7 of the University's Document for -Guidelines and Procedures for Research, March 2012. Name of the Graduate Student

	Signature	Date
Ukula yohannis	_____	_____

Name (s) of the Research Supervisor (s)	Signature	Date
_____	_____	_____

Name of Chairperson	Signature	Date
_____	_____	_____

Name of Internal Examiner	Signature	Date
_____	_____	_____

Name of External Examiner	Signature	Date
_____	_____	_____

ACKNOWLEDGEMENT

Above all loving kindness and faithfulness of the Almighty Allah in bestowing he althea, strength,patients and protection throughout my life. I am very much indebted to my Major Advisor: Dr.Wondemegegh Demise for his constant advice and encouragement during the development of this thesis. I also want to thank my Co-Advisor: Mr. Zewuge Hailu for his constructive comments and guidance of my research work from initial inception up to its completion. I would like to extend my special thanks to the Shashemene City Sport Administration for helping by good idea. From the inner core of my heart, I would like to express my deep sense of gratitude to my families especially my mother Buna Edao and my Husband Dekebo Bulli for their support and immeasurable sacrifices; they paid to reach me at this spot. My sincere appreciation and thanks also go to my instructors of Jimma University Sport Science Department staffs for their constant moral support friendly advice and editorial work the language during the entire research work. A special thank goes to my best Brother Abdisa Yohannis for his unconditional love regular moral and constructive comments of my research work from the initial up to its completion. Finally, I am grateful to thank Jimma University Postgraduate Program Directorate for their cooperation throughout all the necessary support to my works.

TABLE OF CONTENT

ACKNOWLEDGEMENT	I
LIST OF FIGURES.....	VI
ABBREVIATION AND ACRONYMS.....	VII
CHAPTER ONE.....	1
INTRODUCTION	1
1.2. STATEMENT OF THE PROBLEM	3
1.3. RESEARCH QUESTIONS	4
1.4. OBJECTIVES OF STUDY.....	4
1.4.1. GENERAL OBJECTIVE.....	4
1.4.2. SPECIFIC OBJECTIVES.....	4
1.5. SIGNIFICANCES OF THE STUDY	4
1.6 DELIMITATIONS OF THE STUDY	5
1.7. LIMITATIONS OF THE STUDY	5
1.8. OPERATIONAL DEFINITION.....	6
1.9. ORGANIZATION OF THE STUDY	6
CHAPTER TWO.....	7
2. REVIEW OF RELATED LITERATURE	7
2.1. OVERWEIGHT AND OBESITY	7
2.1.1. Body composition	7
2.1.2. Body Mass Index.....	8
2.1.3. Components of Health-Related Physical	9
2.2. HEALTH RELATED PHYSICAL FITNESS TEST	10
2.2.1. Test for Cardiovascular Endurance	10
2.2.2. Test for Muscular Strength.....	10
2.2.3. Test for Muscular Endurance	11
2.2.4. Test for Flexibility.....	11
2.2.5. Test for Body Compositions	11
2.3. PHYSICAL ACTIVITY AND THE PREVENTION OF WEIGHT GAIN	12

2.4. AEROBIC EXERCISE AND ITS BENEFITS	12
2.4.1 Aerobic Exercise	12
2.4.2. Types of Aerobic Exercise	14
2.5. BENEFITS OF AEROBIC EXERCISE.....	14
2.6. AEROBIC EXERCISE INTENSITY ON WEIGHT LOSS.....	15
2.7. BENEFITS OF REGULAR EXERCISE.....	15
2.8. WEIGHT COMPENSATION FOR AEROBIC EXERCISE TRAINING	16
2.9. EFFECT OF AEROBIC EXERCISE ON BODY COMPOSITION	17
2.10. STUDIES ON AEROBIC EXERCISE	17
2.11. WEIGHT LOSS FROM HIGH VOLUME AEROBIC EXERCISE TRAINING WITHOUT CALORIC RESTRICTION	20
2.12. WEIGHT LOSS FROM AEROBIC EXERCISE AT PUBLIC HEALTH RECOMMENDATION LEVELS	20
2.13. EXERCISE AND WEIGHT MAINTENANCE.....	22
2.14. SEDENTARY LIFESTYLE AND ITS EFFECTS	23
2.14.1. Sedentary Lifestyle.....	23
2.14.2. EFFECTS OF SEDENTARY LIFE STYLE.....	23
2.15. CONTRIBUTING EFFECTS OF AEROBIC EXERCISE INTENSITY OR RESISTANCE TRAINING ON	24
CHAPTER THREE.....	26
3. RESEARCH METHODS	26
3.1. STUDY AREA.....	26
3.2. THE STUDY DESIGN	27
3.3. POPULATION OF THE STUDY	28
3.4. SAMPLE SIZE AND SAMPLING TECHNIQUES.....	28
3.5 INCLUSION AND EXCLUSION CRITERIA	29
3.6. SOURCE OF THE DATA	29
3.7 DATA COLLECTION INSTRUMENTS.....	29
3.8 PROCEDURES OF DATA COLLECTION	29
3.9 METHOD OF DATA ANALYSIS.....	32
3.10 ETHICAL ISSUES AND CODE OF CONDUCT	32

CHAPTER FOUR	33
4. RESULTS AND DISCUSSION.....	33
4.1 DISCUSSION	33
4.2 RESULTS OF THE STUDY.....	33
4.2.1 Characteristics of study participant variables.....	33
CHAPTER FIVE	40
5. SUMMARY, CONCLUSION, AND RECOMMENDATION.....	40
5.1. SUMMARY OF THE MAJOR FINDINGS	40
5.2. CONCLUSIONS	41
5.3. RECOMMENDATIONS	42
REFERENCES	43
APPENDICES.....	52

LIST OF TABEL

TABLE1: EXPECTED INITIAL WEIGHT LOSS AND POSSIBLY PRODUCING CLINICALLY SIGNIFICANT WEIGHT LOSS FROM DIFFERENT MODALITIES OF EXERCISE TRAINING.	22
TABLE 2 THE MEAN VALUE OF BODY WEIGHT, HEIGHT, AGE AND BODY MASS INDEX OF BOTH CONTROLS GROUP AND EXPERIMENTAL GROUP.	34
TABLE 6. PAIRED T TEST RESULTS OF WAIST CIRCUMFERENCE.....	37
TABLE 5. PAIRED T TEST RESULTS OF BMI	36
TABLE 4. PAIRED T TEST RESULTS OF BODY WEIGHT	36
TABLE 3.MEAN VALUES OF BODY WEIGHT (KG), WAIST CIRCUMFERENCE (WC)OF (CG) AND (EG); AT DIFFERENT OCCASIONS OF TRAINING PROGRAM.....	35

LIST OF FIGURES

<u>Figure 1.graphical presation of body index mass index</u>	9
<u>FIGURE 2. MAP OF THE STUDY SITE</u>	27

ABBREVIATION AND ACRONYMS

ACSM	American College of Sport Medicine
ANCOVA	Analysis of Covariance
BMR	Basal Metabolic Rate
BMI	Body Mass Index
CG	Control Group
CHD	Coronary Heart Disease
CVD	Cardio Vascular Disease
CSAE	Central Statics Agency of Ethiopia
CSA	Central Statistical Agency
CVE	Cardio Vascular Endurance
DREW	Dose Response to Exercise in Women
EG	Experimental Group
EI	Energy Intake
EIEE	Exercise Induced Energy Expenditure
ET	Exercise Training
HDL	High Density Lipoproteins
HF	Heat Failure
HR _{max}	Maximum Heart Rat
LBM	Lean Body Mass
LDL	Low Density Lipoproteins
MD	Mean Difference
PA	Physical Activities
PT	Pre-test
POT	Post Test
RCTs	Randomized Controlled Trials
SD	Standard Deviation
STRITDE	Studies of a Targeted Risk Reduction Intervention Through Defined Exercise
SPSS	Statistical Package for Social Sciences

US	United States
WC	Waist Circumference
WHO	World Health Organization
WHR	Waist Hip Ratio

ABSTRACT

Introduction: developing effective aerobic exercise program for overweight woman is a strategy for improving BW, BMI and WC. **Objectives:** The general objective of this study is to investigate the effect of 12 weeks aerobic exercise program on the weight loss of sedentary overweight women in Shashemene City. The design was experimental. **Methods:** Pre and post. Based on the standard t-protocol test. The total population's participants are 40 subjects were selected for study and they were assigned into two groups equally using random sampling techniques. Those (n-20) are experimental group and (n-20) control group. The data collection instrument was quantitatively participated in the typical training of 40-60 per week but only EG participated 3 sessions per week which comprised as situp, walking, jogging, running and rope skipping. **Results:** This indicates that moderate-intensity aerobic exercise training program were effective for the reduction of body weight and waist circumference for overweight and obese women. Sedentary female's community may be encouraged to undergo this type of exercise training regularly for physical fitness improvement. 40 overweight women whose BMI was 26.52 – 24.75 kg/m² and age in between 25-35 years participated in this study. Pre and Post tests were taken and analyzed accordingly. The mean values of participants' age were 30.15 for control group and 30.15 experimental group respectively. The mean values of participants' height were 1.65 for control group and 1.65 for experimental group respectively. The mean value of body weight before training was 72.15 for control and the mean value of body weight after training was 73.25 for control group. The mean value of body weight before training was 71.20 and after training 67.15 for experimental group. The mean of body weight was increased by -1.1 for control group and 4.05 was reduced in experimental group throughout the study period. The participants were reduced significantly due to aerobic training. In waist circumference test, the mean value WC before training was 90.80 and after training was 91.75 for control. Before training 91.45 and after training 82.90 for experimental group, control group was increased by -0.95 but in experimental group waist circumference was decreased by 8.55 markedly. The data collection study subject was analyzed using SPSS version 20 software by paired t-test with level of significant less than 0.05 considered. The result showed aerobic exercise significantly improved Body weight, Body mass Indices and waist circumference in EG ($P < 0.05$). Besides, there were significant differences were found in all of the variables in CG ($P > 0.05$). **Conclusion:** Based on the findings, it was concluded that moderate-intensity aerobic exercise program had a positive effect on body weight reduction and waist circumference reduction of the subjects.

KeyWords: Aerobic training, BMI, body weight, Sedentary women, Waist Circumference

CHAPTER ONE

INTRODUCTION

1.1 BACKGROUND OF THE STUDY

Overweight and obesity are a major public health problem worldwide. For example, the World Health Organization reported that the global prevalence of overweight and obesity in adults, defined as a body mass index (BMI) ≥ 25 kg/m² is approximately 2 billion and by the year 2015 will increase to approximately 3 billion (George *et al.* 2012). Obesity is a major public health problem in the world. The number of obese peoples increases significantly and developing countries are highly concerned (World Health Organization, 2010).

Overweight is associated with long-term health and reduced life quality. It has been recommended that effective weight loss strategies be developed. Although caloric restriction has been the major weight loss strategy, it has been shown that exercise programs designed for fat loss result in an increase in cardio respiratory fitness and a preservation of fat-free mass. Most exercise programs designed for weight loss have focused on steady-state exercise (SSE) of around 30min at a moderate intensity on most days of the week. Disappointingly, these kinds of exercise programs have led to little or no fat loss. Thus, what is needed is an exercise protocol that can be carried out by overweight, inactive individuals that more effectively induces fat loss. The increase of exercise capacities and physical fitness result in several health benefits (Vendees *et al.*, 2007).

Aerobic-type training of between 225 and 420 min/week is recommended to those who wish to lose fat mass by increasing physical activity levels (Swift *et al.*, 2014) but very high intensity, short duration training might also be effective (Trapp *et al.*, 2008). The Australian National Physical Activity Guidelines for Adults recommend more than 30 min of daily physical activity (William *et al.*, 2016). Numerous studies have investigated the effects of exercise training, demonstrating significant improvements to CVD risk factors after aerobic exercise training. However, it is unclear whether health benefits are limited to aerobic training or if other exercise modalities such as resistance training or a combination are as effective or more effective in the overweight and obese (Sullen *et al.*, 2012).

An excess of energy intake relative to energy expenditure is the central reason for the development of obesity. It occurs due to storage of excess energy as adipose tissue. As the

incidence of obesity increases, the significant health and economic consequences also increase. Obesity, which has been linked to a variety of chronic diseases results in almost 300000 deaths each year and causes 117 billion dollars worth of direct and indirect annual costs in the United States.(A.Yes *et al.*,2006). Aerobic exercise is a commonly prescribed method of weight management with sustained increases in daily energy expenditure intended to promote energy deficit and weight loss. However, the regulation of energy balance is a dynamic process in which individual components of energy balance interact in a coordinated fashion. Consequently, adjustments in exercise induced energy expenditure (EIEE) may elicit compensatory changes in other components of energy balance that attenuate the prescribed energy deficit and subsequent weight loss. Indeed, compensatory changes in hunger and energy intake (EI) are commonly cited as reasons why exercise often produces modest weight loss (1.5 to 3.0 kg over 3 to 12 months) that is less than expected theoretically (Thomas *et al.* 2012).

Obesity is a major risk factor for many cardiovascular (CV) diseases such as coronary heart disease (CHD), heart failure (HF), stroke, ventricular dysfunction and cardiac arrhythmias (Damos *et al.* 2014). Overweight and obesity is fundamentally caused by energy imbalance between calories consumed and calories expended (Musa *et al.* 2013). Aerobic training (AT) comprises several modes of activities that primarily stress the aerobic energy system and produce a number of cardiovascular (CV) and respiratory adaptations that increase endurance. High levels of aerobic fitness are mandatory for endurance athletes such as cyclists, distance runners, athletes and swimmers (Ratamess, 2012).

In the United States (US) an estimated 68% of adults are either overweight or obese. Fat accumulation and especially visceral fat (i.e. abdominal fat that surrounds the vital organs on the trunk and stomach area of the body) is associated with cardiovascular diseases type II diabetes, metabolic syndrome, excess weight, weakness, falls and fatigue (Joav *et al.* 2013).During exercise substrate metabolism primarily depends on exercise intensity, such that absolute rates of fat oxidation increase and then decline with increasing intensity, whereas rates of carbohydrate oxidation progressively increases with exercise intensity. This is known as the “crossover concept” of exercise substrate metabolism. The dynamics of fat oxidation with increasing exercise intensity can be depicted as a bell-shaped relationship (Pernille *et al*

2015). The prevalence of overweight and obesity has increased to epidemic proportions in the industrialized world and it is now dramatically rising in low and middle income countries particularly in urban settings. It is well known that regular physical activity (PA) provides health benefits and it is considered an essential component of primary and secondary prevention form of metabolic- syndrome related pathologies. Recent experimental data suggests that subjects who increased their level of PA over time have a decreased mortality rate compared to those who were consistently unfit (Gianni *et al.*, 2015).

1.2. Statement of the Problem

As the topic is “Effects of 12 week aerobic exercise program on the weight loss of sedentary overweight selected variables in the case of sedentary over weight woman in Shashemene city from female Assosecion namely Eder. This research was conducted to show that scheduled aerobic exercise programs for female Assosecion the name of Eder in Shashemene city may lead to a decrease in body weight. Waist Circumference and BMI the capacity of aerobic exercise. Because of sedentary life a lot of women were attacked by chronic disease as a result of overweight obesity. Aerobic exercise is essential components that seriously addresses and improve the three variable of woman Assosecion. Many studies believe that aerobic exercise can have miraculous health benefits on cardio respiratory and body composition improvement. However, it is expected that after three month the selecting woman will recognize the importance and in aerobic exercise designed to maintain and enhance healthy lifestyles (Connecticut State Department of Education 2009).

Many research studies shows the effect of different exercises for the development of physical fitness but now a day’s few studies were done specifically on the area of aerobic exercise training on selected physical fitness variables. Aerobic exercises are one of important physical activities which are at center of attention in recent years (Yildirim, 2012). Today, most of women aware of aerobic exercise can positively affect physical fitness. Physical fitness training is mandatory for every individual to maintain good health and to prevent body weight. Aerobic exercises is more vital to burn out extra calories that stored in human`s body. So aerobic exercise is very important to improve the excess weight of the body aerobic capacity (cardiovascular endurance) as well as reduce the chance of being exposed to chronic diseases.

1.3. Research questions

Therefore, this study will be tried to answer the following research questions: -

1. What is the effect of aerobic exercise on the bodyweight of sedentary women?
2. What is the effect of aerobics exercise on the waist circumference of sedentary women?
3. Does the aerobics exercise program make any change in the body mass index of sedentary overweight women?

1.4. Objectives of Study

1.4.1. General Objective

- The general objective of this study is to examine the effect of 12 weeks of aerobic exercise program on weight loss of sedentary overweight women in Shashemene City.

1.4.2. Specific Objectives

1. To examine the effect of aerobic exercise on improvement of body weight of sedentary Overweight women.
2. To assess the effect of moderate-intensity aerobics exercise on the waist circumference of sedentary women?
3. To identify the effect of aerobics exercise program on the body mass index of sedentary overweight women?

1.5. Significances of the Study

Currently, enhancement of body weight efficiency is designed upon critical study of human physiology, modern way of feeding and scientific way of training based upon a new findings and principles of investigation.

- ✓ This research will contribute in addressing the effects of 12weeks aerobics exercise program for overweight women and managers to understand formulate and implement on designing effective strategies for aerobics exercise program.

- ✓ The study also helps to develop our country's overweight and obese individuals to become health and physically fit as of the world class by improving their physical fitness.
- ✓ In addition, it will help for others as a research work for depth studies on the problem undertaken.
- ✓ The finding of the research may help as reference for researchers who will conduct advanced researches of exercise physiology particularly in aerobics exercise program effect.

1.6 Delimitations of the study

In research, delimitations address how the study is narrowed in scope (Creswell, 1998). The study was designed to investigate the effects of twelve(12) week aerobic exercise program on overweight women in Shashemene city. This study was delimited in the following areas.

- ❖ Subjects were selected at Shashemene city women Assosecion namely Eder, age ranged between 25-35 years. And healthy untrained that has not any physical disabilities or medical conditions and volunteers participated in this study.
- ❖ Selected Bodyweight variables were cardio respiratory fitness measured using 12 minute run test and step test and body composition measured using body mass Idex.
- ❖ Aerobic exercise trainings were running, cross running, walking, jogging and rope skipping exercises.
- ❖ The time of training was limited to three days per week and 60minutes per sessions. The study were conducted in the academic year of 2009 E.C/2016-2017 G.C
- ❖ The time of training was limited to three days per week and 60minutes per sessions.

1.7. Limitations of the Study

In this research, It is difficult and unmanageable to conduct study all sedentary overweight women in Shashemene city. Therefore, the number of participant are limited to 40 subject only and follow up were done on one area namely Shashemene high school which is found in Shashemene city.

1.8. Operational Definition

Aerobic Exercise any activity that uses large muscle groups can be maintained continuously and is rhythmic in nature and it is a type of exercise that overloads the heart and lungs and causes them to work harder than at rest ((Kraus *et al.*,2002).

Obese abnormal accumulation of body fat usually, 20% or more over an individual's ideal body weight (Mills and Mae 1994).

Overweight is abnormal or excessive fat accumulation that may impair health.

Sedentary Women women doing a lot of sitting not doing or involving much physical activity.

1.9. Organization of the study

The study is organized and presented in Five chapters. The first chapter gives a general idea on the background of the study, statement of the problem, explain the significance of the study and describe briefly the scope of the study. The second chapter is Review Literature and The Third one is Materials and Methods. The fourth chapter is Results and Discussion. The last and fifth is Summary, Conclusion and Recommendation.

CHAPTER TWO

2. REVIEW OF RELATED LITERATURE

This chapter includes overweight and obesity body composition components of health related physical fitness health related physical fitness test for cardiovascular endurance test for muscular strength test for muscular endurance test for flexibility test for body compositions physical activity and the prevention of weight gain aerobic exercise and its benefits aerobic exercise. Types of aerobic exercise benefits of aerobic exercise Aerobic exercise intensity on weight loss, benefits of regular exercise weight compensation for aerobic exercise training effect of aerobic exercise on body composition studies on aerobic exercise weight loss from high volume aerobic exercise training without caloric restriction, weight loss from aerobic exercise at public health recommendation levels Exercise and weight maintenance sedentary lifestyle and its effects sedentary lifestyle contributing effects of aerobic exercise intensity or resistance training on weight loss.

2.1. Overweight and Obesity

Although being overweight is what bothers most people it is really the amount and location of fat (%BF, abdominal fat mass) that should be of concern. Excess weight can be caused by high levels of lean muscle mass but additional muscle mass is beneficial. Except in rare instances such as providing protection from the cold water for an English Channel swimmer or certain wasting diseases excess fat is generally not beneficial. There are no universally agreed upon acceptable %BF standards. The most typically used normal values for young adults (20–29 yr) are 12–15% for males and 22–25% for females with an allowance of an additional 2% for each decade of age. Obesity is defined as +5% BF above the normal value (Kaminsky and Dwyer2006).

2.1.1. Body composition

Body composition is defined as the partitioning of body mass in to FFM (weight or percentage) and fat mass (weight or percentage). Compartmentalizing the body in to only fat and FFW (not water, mineral, protein, and fat) and using this two-compartment model to determine percent body fat (%BF) depends on the following assumptions:-

- ✚ The densities of the fat and the FFW are known and additive.
- ✚ The densities of water, bone mineral, and protein that make up the FFW are known and are relatively constant from individual to individual.
- ✚ The percentage of each fat-free component is relatively stable from individual to individual.
- ✚ The individual being evaluated differs from the assumptions of the equation being used only in the amount of storage fat.


Skin folds

The most widely used anthropometric estimation of body size or composition involves the measurement of skin folds at selected sites. Skin folds (sometimes called fat folds) are the double thickness of the skin plus the adipose tissue between the parallel layers of the skin (Figure 7.4). Because skin thickness varies only slightly among individuals skin fold measures generally indicate the thickness of the subcutaneous fat (Behnke and Wilmore, 1974). Technically however adipose tissue (and thus the subcutaneous fat fold) has both a fat component and a fat-free component. The fat-free component is composed of water blood vessels and nerves. As the fat content of the adipose tissue increases (as in obesity) the water content decreases (Roche, 1987). The use of skin fold thicknesses to estimate body composition is based on two assumptions. The first is that selected skin fold sites are representatives of the total subcutaneous adipose tissue mass. In general evidence supports this assumption (Roche, 1987). The second assumption is that the subcutaneous tissue mass has a known relationship with total body fat. Table 7.2 shows the distribution of total body fat and the relative percentages of each storage site for a reference male and a reference female 25–35y ears old.

2.1.2. Body Mass Index

Mass Index (BMI) is a ratio of the total body weight to height. Several ratios have been proposed but the one used most frequently is weight (in kilograms) divided by height (in meters) squared [$WT \div HT^2$ ($kg \cdot m^2$)]. This ratio is also known as the Quenelle index (Boride, 1988). Calculated BMI can then be compared against standard values to determine whether the individual has acceptable body weight is overweight or is obese.

Figure 1.graphical presation of body index mass index



Body Mass Index(BMI)

Source;<https://www.google.com.et/search>

2.1.3. Components of Health-Related Physical Fitness

Physical fitness is the ability to function effectively in physical work training and other activities and still have enough energy left over to handle any emergencies which may arise. The components of health-related physical fitness are as follows:

Cardiovascular Endurance: the efficiency with which the body delivers oxygen and nutrients needed for muscular activity and transports waste products from the cells. Cardiovascular endurance sometimes called cardio respiratory fitness aerobic fitness or aerobic capacity is one of the basic components of physical fitness. Cardio respiratory fitness is a condition in which the body's cardiovascular (circulatory) and respiratory systems function together especially during exercise or work to ensure that adequate oxygen is supplied to the working muscles to produce energy. Cardio respiratory fitness is needed for prolonged rhythmic use of the body's large muscle groups. A high level of cardio respiratory fitness permits continuous physical activity without a decline in performance and allows for rapid recovery following fatiguing physical activity.

Muscular Strength: refers to the force or tension that can be generated by a muscle or muscle group during one maximal effort (Physiology of Exercise: Responses and Adaptations, 2nd edition).

Muscular Endurance: the ability of a muscle or muscle group to perform repeated movements with a sub-maximal load for extended periods of times (Gut in, 1980).

Flexibility: Flexibility is the degree to which body segments can move or be moved around a Joint. (Brown, 1986)

2.2. Health Related Physical Fitness Test

2.2.1. Test for Cardiovascular Endurance

Twelve-minute run / walk test: This test objective is to measure the cardiovascular endurance of the participants. For this test the participants will run for 12 minutes, and the total distance covered will be recorded. The participants can walk also though the participants will be encouraged to push's them as hard as they could. The average distance for men is 2200-2399m for the age 20-29, 1900-2299m for the age 30-39, 1700-2099m for the age 40-49, 1600-1999m for the age 50. For the female 1800-2199m for the age 20-29, 1700-1999m for the age 30-39, 1500-1899m for the age of 40-49, 1400-1699m for the age 50 (Cooper, 1968).

2.2.2. Test for Muscular Strength

Push up test: The push-up test is a basic fitness test used by coaches trainers and athletes to assess upper body fitness and to monitor progress during strength and fitness training. This test helps you compare your own upper body muscular endurance to others of your age and gender and track your fitness program over time. Men should use the standard "military style" pushup position with only the hands and the toes touching the floor. Women have the additional option of using the "bent knee" position. To do this, kneel on the floor, hands on either side of the chest and keep you back straight. The average number of pushups for men is 19-34 for the age 17-19, for the age 20-29, 13-24 for the age 30-39, 11-20 for the age 40-49, 9-17 for the age 50-59, 6-16 for the age 60-65 and for women's is 11-20 for the age 17-19, 12-22 for the age 20-29, 10-21 for the age 30-39, 8-17 for the age 40-49, 7-14 for the age 50-59, 5-12 for the age 60-65 (Golding *et al.*, 1986).

2.2.3. Test for Muscular Endurance

Sit up Test: The objective of this test is to measure abdominal muscular strength and endurance of the abdominals and hip-flexors important in back support and core stability. For this test Subjects lays on her back on the mat with knee bent and feet about two feet apart. Her hands are placed on the back of the neck with the fingers interlocked elbows are retracted. Assistance data collector holds the subject ankles down the heels being in contact with the mat at all times. The subject sits up, turning the trunk to the left and touching the right elbow to the left knee returns to starting position and then sits up turning the trunk to the right and touching the left elbow to the right knee. The maximum number of sit ups done within 30 seconds will be taken as her score. The average sit ups for female is 29-39 for the age 18-25, 25-28 for the age 26-35, 19-22 for the age 36-45,14-17 for the age 46-55,10-12 for the age 56-65, 6-10 for the age 65+ (Golding *e al.*1986).

2.2.4. Test for Flexibility

Sit and reach test: This test measures the flexibility of the lower back and hamstring muscles. It involves sitting on the floor with legs out straight ahead. Feet shoes ohms are placed with the soles flat ageist the box shoulder width apart. Both knees are held flat against the sit and reach apparatus by the tester. With hands on top of each other and palms facing down the subject reaches forward along the measuring line as far as possible. After three practice reaches the fourth reach is held for at least two seconds while the distance is recorded. Make sure there is no jerky movements and that the fingertips remain level and the legs flat. The score is recorded to the nearest centimeter as the distance before (negative) or beyond (positive) the toes. The average score for boys is between +0 to +5cm and girls is +1 to +10cm (Wells and Dillon,1952) .

2.2.5. Test for Body Compositions

Body mass index: BMI (Body Mass Index) is a measurement of body fat based on height and weight that applies to both men and women between the ages of 18 and 65 years. BMI can be used to indicate if you are overweight obese underweight or normal. A healthy BMI score is between 20 and 25. A score below 20 indicates that you may be underweight; a value above 25 indicates that you may be overweight (www.bmi-calculator.net). BMI is just a guide it does

not accurately apply to elderly populations, pregnant women or very muscular athletes such as weight lifters (WHO, 2012).

2.3. Physical Activity and the Prevention of Weight Gain

Changes in weight are affected by the amount of energy expended versus the amount of energy consumed (Thom set *et al.* 2012). Therefore, if the energy expenditure remains low but dietary consumption levels are in excess weight gain will occur. Several researchers have argued that declines in physical activities in occupational and leisure settings (Fogelholm, 2000) may have an important role in the increase in obesity rates over the last 30+ years. Furthermore, many epidemiological studies suggest that physical activities have an important role in weight gain (Saris *et al.* 2003).

(Williams *et al.* 2007) using data from the National Health and Nutrition Examination Survey observed that low levels of self-reported recreational physical activities is associated with a 3-fold greater risk of major weight gain in men and almost a 4-fold in women. In a prospective study of 34,079 middle aged women (mean: 52.2 years). (Pulse *et al.*,2010). Observed that in women physical activities Author Manuscript in less than 7.5 metabolic equivalent hr./wk. compared to women who participated in greater than (Saris *et al.*, 2003) metabolic equivalent hr./wk. (approximately 300 minutes/week of moderate physical activities).

Several studies using the aerobic center longitudinal study database have observed that CRF level (Van *et al.*, 2007) physical activities level and change in physical activities level 24 are inversely associated with future weight gain. Clearly physical activities and CRF levels have an important role in weight gain for those at risk.

2.4. Aerobic Exercise and its Benefits

2.4 .1Aerobic Exercise

Aerobic exercise is physical exercise of relatively low intensity that depends primarily on the aerobic energy-generating process. Aerobic literally means “living in air” and refers to the use of oxygen to adequately meet energy demands during exercise via aerobic metabolism. Generally light-to-moderate intensity activities that are sufficiently supported by aerobic metabolism can be performed for extended periods of time. Aerobic exercises are a wonderful

way to burn your fat and tone your body muscles leaving you healthy and in a good shape. Finding the perfect Workout Routines takes time and effort. These best workout routines is a great place to start if a person is interested in flat abs. All these activities are healthy easier to perform and inexpensive. Aerobic exercises are beneficial in so many ways like strengthening the respiratory muscles strengthening and enlarge the heart muscle and improve its pumping, improving blood circulation and red blood cells reducing stress and depression increasing your stamina and endurance of your muscles. In short it reduces the risk of heart attacks. (International Journal of Physical Education, Sports and Health, 2015). Aerobic exercise is any activity that uses large muscle groups, can be maintained continuously for a long period of time and is rhythmic in nature. Aerobic activity trains the heart, lungs and cardiovascular system to process and deliver oxygen more quickly and efficiently to every part of the body. As the heart muscle becomes stronger and more efficient, a larger amount of blood can be pumped with each stroke. Fewer strokes are then required to rapidly transport oxygen to all parts of the body. An aerobically fit individual can work long more vigorously and achieve a quicker recovery at the end of the aerobic session. Regardless of age, weight or athletic ability aerobic activity is good for human being. As the body adapts to regular aerobic exercise, we will get stronger and fitter (American College of Sports Medicine, 2010).

According to Braun, et al (2013) and DHHS (2013) aerobic exercise can help to feel better and enjoy life to the fullest in the following way: -

Keep excess pounds at bay: - Combined with a healthy diet, aerobic exercise helps to lose weight and keep it off.

Increase stamina: - aerobic exercise may make tired in the short term. But over the long term, you will enjoy increased stamina and reduced fatigue.

Ward off viral illnesses: - Aerobic exercise may activate immune system. This may leave less susceptible to minor viral illnesses such as colds and flu.

Reduce health risks: - Aerobic exercise reduces the risk of many conditions, including obesity heart disease high blood pressure, type two diabetes metabolic syndrome stroke and certain types of cancer. Weight bearing aerobic exercises, such as walking reduces the risk of osteoporosis.

Stay active and independent as you age: - Aerobic exercise keeps muscles strong, which can help to maintain mobility as you get older.

Strengthen heart: - A stronger heart doesn't need to beat as fast. A stronger heart also pumps blood more efficiently which improves blood flow to all parts.

2.4.2. Types of Aerobic Exercise

Any activity that uses large muscle groups can be maintained continuously and is rhythmical in nature can be regarded as an aerobic exercise. In general aerobic exercises requiring little skill to perform are more commonly recommended for all adults to improve fitness. Aerobic exercises that require minimal skills and can be easily modified to accommodate individual physical fitness levels include brisk walking, leisure cycling, swimming, and aqua-aerobics and slow dancing. Aerobic exercises that are typically performed at a higher intensity and therefore, are recommended for persons who exercise regularly include jogging, running, aerobics, stepping exercise fast dancing and elliptical exercise. These exercises should be performed with the recommended dosage of aerobic exercise which means (frequency (F) intensity (I) and duration (time, T) of the exercise performed. In combination with the type (T) of exercise performed these factors constitute the basic components of the core principle of exercise prescription (the FITT principle). It should be noted that even small increases in caloric expenditure with physical activity may improve physical fitness out comes (General principles of exercise prescription 4thed).

2.5. Benefits of Aerobic Exercise

Lifestyle plays an essential part in maintaining long-term cardiovascular health. A healthy diet moderate drinking, not smoking and plenty of exercise can all help to maintain a healthy cardiovascular system. Inactivity is one of the major risk factors for heart disease and exercise helps improve heart health and can even reverse some heart disease risk factors. Dr. Kelly & Associates 1989 noted that aerobic exercise produces beneficial effect on human bodies by strengthening the muscles involved in respiration, improving circulation efficiency and reducing blood pressure, reducing body fat, reduced stress and preventing health problems.

In addition, aerobic exercise has various effects on human body including, reduced cholesterol and blood pressure, improved muscular endurance reduced body fat, increased metabolism strengthen heart and lungs improved quality and quantity of life. Exercise not only extends your life, but also gives you more energy to live it to the fullest. Aerobic exercise improves the

strength of bones, ligaments and tendons allows body to use fats and sugars more efficiently, burn lots of calories and plays an important role in reducing the onset and symptoms of aging and illness. Aerobic exercise reduces risk of heart disease, vascular disease and diabetes and can help those trying to quit smoking by relieving cravings and improving lung function (Kathleen, 2006).

2.6. Aerobic Exercise Intensity on Weight Loss

Vigorous intensity aerobic has been shown to have enhanced health benefits for important risk factors including visceral fat measures of glucose/insulin metabolism (Dippier *et al.*, 2006) compared to moderate intensity aerobic training. In terms of weight change, when different intensities of aerobic exercise training are matched for caloric expenditure or aerobic training dose, both vigorous and moderate intensity aerobic exercise training result in similar amounts of weight loss. (O'Donovan *et al.* 2005) in resistance training of 64 obese men observed similar changes in weight following 6 months of moderate intensity (-1.1 kg) and high intensity (-0.5 kg) aerobic exercise training. In the study of (Kraus *et al.*,2002) observed similar weight loss in overweight/obese adults participating in 8 months of moderate (-0.6 kg) and high intensity (-0.2 kg) aerobic exercise training at the same exercise dose (14 kilocalories per kg per week). Vigorous intensity aerobic exercise training can contribute to greater weight loss if matched for session time compared to moderate intensity aerobic exercise training because the total energy expenditure is greater. Thus, if vigorous aerobic exercise training can be sustained and is enjoyable for the patient, the exercise program may induce additional health benefits and potentially increase the energy expenditure seen with exercise training (if replacing the exercise time of moderate intensity training).

2.7. Benefits of Regular Exercise

Exercise mode recommendations for specific health benefits remain unclear due to in large part to the sparse scientific data supporting this recommendation. Approximately two-thirds of United States adults are overweight or obese. Although professional organizations have historically focused exercise guidelines on endurance or aerobic training(at) for weight loss and maintenance (Jakicic *et al.*,2001). recent guidelines and position statements targeting body weight reduction and maintenance have suggested that resistance training may also be

effective for reducing fat mass. In some cases guidelines may lead to misperceptions among clinicians, exercise professionals, and laypersons about the strength of the evidence regarding the effectiveness of resistance training for inducing weight and fat mass loss (Williams *et al.* 2007) leading the reader to believe that resistance training has been conclusively shown to reduce fat mass. However, close examinations of the published literature reveals that randomized controlled trials are inconclusive on this point (Castaneda *et al.*, 2002).

2.8. Weight Compensation for Aerobic Exercise Training

Individuals who lose less weight than expected based on their training energy expenditure have been termed “weight compensators.” Several studies have examined weight compensation after aerobic exercise training (King *et al.*, 2008). observed increased energy intake and increased fat intake in weight compensators compared to those that did not compensate for weight loss.

Using data from the (Church *et al.*,2009) examined weight compensation in postmenopausal women who were required to perform exercise training at 50%, 100% and 150% of public health recommendations and restricted the analysis to those who were 85% compliant to exercise training. The authors observed that the most weight compensation (less weight loss achieved than predicted from exercise training alone) occurred in the women exercising at 150% of the recommended volume. In fact, the amount of actual weight loss achieved in women exercising at 50% (-1.4 kg) and 150% (-1.5 kg) of the physical activity’s recommendations are virtually identical to each other despite the greater level of energy weekly energy expenditure in the 150% groups.

(Thomas *et al.* 2012) performed an analysis of weight change from aerobic exercise training interventions and concluded that the major factors limiting the expected weight loss from aerobic exercise training were dietary compensation and low aerobic exercise training dose. At the present time evidence is limited to explain whether other factors of the energy balance equation, including compensatory changes in non-exercise physical activities (except for perhaps older adults) resting metabolic rate, movement efficiency or changes in lean mass are responsible for weight compensation with exercise training

2.9. Effect of Aerobic Exercise on Body Composition

Regular aerobic exercises reduce the body fat percentage without the loss of the muscle as well as important effect on anthropometric and hematologic level of obese and overweight women (Ervine *al.* 2010). Aerobic exercise including walking, running, and swimming has been proven to be an effective way to lose weight. Body Composition Changes for over fat or borderline over fat people, regular aerobic exercise reduces body mass and body fat. Increases in fat free body mass also accompany a regular program of resistance training. Exercise only, or exercise combined with calorie restriction reduces body fat more than fat lost with only dieting because exercise conserves the body's lean tissue mass (Katch *et al.*,2011). Cardiovascular endurance (CVE) is one of the most important measures of overall health.

A person's level of cardiovascular endurance helps predict probability of disease, quality of life, and ability to react to acute physical and mental stress. For healthy individuals, higher cardiovascular endurance also indicates an elevated level of physical fitness (Corbett, 2009).

2.10. Studies on Aerobic Exercise

(Selvam and Sudha2008) conducted a study on selected effect of aerobic exercise on selected physiological variables among college girls. For this study aerobic exercise uses large muscle groups rhythmically and continuously and elevates the heart rate and breathing for a sustained period. Common examples include walking, jogging/running, swimming, rowing, stair climbing, bicycling, cross country skiing, step and dance exercise classes, roller skating, and the more continuous forms of tennis, racquet ball and squash. To achieve this purpose, 60 girls were selected from The ivannai Amal College for women, Villupuram. The age group of the subjects ranged between 18 to 20 years. The selected subjects were divided into two groups. The groups first trained for aerobic exercise. The training group underwent the training for 5 days in a week for eight weeks and group second acted as control group to make adjustments for differences in the initial means and test the adjusted post-test means for significant differences. The researcher used analysis of covariance (ANCOVA) for interpreting the results. The results for the study revealed that aerobic exercise had a significant effect in the improvement of the physiological variables such as resting pulse rate, breath holding time, vital capacity and respiratory rate (Selvalakshmi 2007). Conducted a study on the effect of

varied aerobic training programs on obese women working in IT companies for the purpose of the study. For this study, the obese women were grouped into three namely, control, floor aerobic and step aerobics group. The collected data on the cardio respiratory parameters prior to and after 12 weeks of varied aerobics training were statistically analysed using analysis of covariance (ANCOVA) and result on vital capacity showed significant improvement due to varied aerobic exercises, as where no significant improvement was found in resting heart rate.

(Oscan and Oz Turk , 2011) in Mug ale, Turkey conducted the study on the effect of twelve week aerobic exercise programme on health related physical fitness components and blood lipids in obese girls .The aim of the study was to investigate the effects of 12 week aerobic exercise program on health related fitness components and blood lipids in obese girls. In this study, a total of 40 girls were recruited as exercise group (n = 20) and control group (n = 19). Participants joined sessions for 60 min per day, 3 days per week for 12-week. There were significant differences in weight, body mass index (BMI), flexibility, sit-ups, hand grip for both hands , skin fold measurements (thigh, triceps, biceps, abdomen, super iliac, sub scapula, chest, body fat percent, heart rate, high density lipoproteins (HDL) low density lipoproteins (LDL) total cholesterol and triglyceride between pre-test and post test scores in the exercise group ($p < 0.05$). It was concluded that regular aerobic exercise may affect health related fitness components and blood lipids positively in girls. Furthermore, it may result in decreasing obesity in girls(Mills and Mae 1994). conducted a study on the effect of low intensity aerobic exercise on muscle strength, flexibility and change of balance among sedentary elderly person. The purpose of this study was to determine the effects of a low intensity aerobic exercise program on muscle strength and flexibility of the lower extremities and balance among sedentary elderly persons. This pre and post-test quasi-experimental study consisted of 47 sedentary elderly subjects not engaged in regular exercise and living in metropolitan housing in South Western Ohio. Convenience sampling was used with two apartment complexes randomly assigned to the experimental or comparison groups. To prevent diffusion of treatment subjects were assigned to these groups depending on their place of residence The 20 experimental subjects, with a mean age of 75.3 participated in eight weeks with low intensity of aerobic exercise while the comparison group (n=20) with a mean age of 74.8, maintained their usual level of activity for eight weeks.

Experimental subjects also did the exercise on their own between classes. The exercise group had significantly greater flexibility of the ankles and knee than the comparison group. No significant differences were found between the groups for muscle strength. Although balance and perception of balance were not significantly different between the groups; the experimental group improved their balance by 22.4% from pre-test.

(Arslan, 2011) conducted the study on the effects of an eight-week step-aerobic dance exercise programme on body composition parameters in middle-aged sedentary obese women in Astray Turkey . This study comprised an eight-week randomized controlled trial. For this study a total of 49 healthy sedentary obese women participated voluntarily. They were randomly divided into two groups: those undertaking a step-aerobic dance exercise programme (n=29) and a control group (n=20). The subjects took part in a step-aerobic dance exercise programme for one hour per day three days a week for eight weeks. The subjects' Body Mass Index (BMI)body weight, waist circumference, waist-hip ratio, four-site skin fold thickness, fat percentage, basal metabolic rate and lean body mass were assessed before and after the completion of the step-aerobic dance exercise programme. After the eight weeks of the step-aerobic dance exercise programme, significant differences were found in the subjects' weight, BMI, body composition parameters, waist-hip ratio (WHR), waist circumference (WC), fat percentage, lean body mass (LBM) and basal metabolic rate (BMR) in the experimental group ($p < 0.05$). There were no significant differences in the control group after the experiment in terms of the same measures ($P > 0.05$) .The result of this study concluded that the step aerobic dance programme proved to be a useful exercise modality for weight loss and in terms of body composition. There was a clear response to the eight-week step aerobic dance programme in terms of central obesity in sedentary obese Turkish women.

2.11. Weight Loss from High Volume Aerobic Exercise Training without Caloric Restriction

Current ACSM recommendations state that exercise programs need to exceed 225 min/wk. to possibly induce clinically significant weight loss(Donnelly *et al.*2009). Supervised ET studies which have demonstrated clinically significant weight loss with aerobic ET (without caloric restriction) have far exceeded the minimum levels of physical activity according to public health definitions(Ross *et al.*, 2000). observed an 8% weight loss in obese men after 12 weeks of aerobic ET with no alterations in dietary habits (daily exercise sessions of 700 kcal). In a different study (Ross *et al.*, 2004) observed a 6.8% weight loss in premenopausal women (BMI > 27) following 14 weeks of aerobic ET with an energy expenditure of 500 kcal per session. In the Midwest Exercise Trial, Donnelly, (n = 131) observed a 5.3% weight loss in men after 16 weeks of aerobic ET at approximately 2,000 kcal per week. In contrast, the women in the exercise group did not have a significant change in weight (0.7 kg) following the intervention, but the exercise program prevented the weight gain observed in the control group (2.9 kg). Thus, clinically significant weight losses are possible with aerobic ET without caloric restriction but it require a high ET volume. For the general population, these ET volumes may not be practical or sustainable.

2.12. Weight Loss from Aerobic Exercise at Public Health Recommendation Levels

Clinical trials of ET that report no weight loss or modest weight loss (<5 kg) still report numerous health benefits for overweight and obese adults with risk factors for disease. These benefits include improving CRF (Johannsen *et al.*, 2013).Weight loss as a result of aerobic ET is very heterogonous and the overall response is related not only to total energy expenditure, but also compensatory changes in dietary caloric intake(Thomas *et al.*, 2012). Large randomized controlled trials (RCTs) which have evaluated the change in weight following aerobic ET programs consistent with PA recommendations have observed either no changes in weight or only modest weight loss. The following studies represent the strongest research design to evaluate changes in weight from aerobic ET as they have a large sample size of overweight or obese individuals at baseline supervised ET sessions, strong adherence to their

aerobic ET program and comparison of weight change against a control group. The Dose Response to Exercise in Women (DREW) study (Church *et al.*, 2007) (n = 464) observed no significant changes in body weight in postmenopausal women exercising at 50% (-0.4 kg), 100% (-2.2 kg) and 150% (-0.6 kg) of public health guidelines for 6 months despite greater than 89% adherence in all ET groups. The Inflammation and Exercise study (n = 129) (Church *et al.*, 2010) observed no significant change in body weight (-0.4 kg) compared to the control group (0.1 kg) after 4 months of ET in adults with elevated C-reactive protein levels at baseline. The Studies of a Targeted Risk Reduction Intervention through Defined Exercise (STRRIDE) study (Kraus *et al.*, 2002) (n = 84) observed significant, but minimal weight loss in those exercising at low amount/moderate intensity (-0.6 kg, 176 min/wk). low amount/high intensity (-0.2 kg, 117 min/w). Or high amount high intensity (-1.5 kg, 171 min/wk.) following 6 months of aerobic ET. The Diabetes Aerobic and Resistance Exercise (DARE) study (Sigel *et al.*, 2007) (n = 251) observed significant weight loss in the aerobic ET group (-0.74 kg) compared to the control group after 22 weeks of intervention in adults with T2DM. Thus, overweight and obese adults who adhere to an exercise program consistent with public health recommendations without a dietary plan involving caloric restriction can expect to experience weight loss in a range of no weight loss to approximately 2 kg. Clinicians should caution their patients that the chances of substantial weight loss are unlikely at these ET levels without caloric restriction. (Donnell *et al.*, 2009) Regardless of the amount of weight loss, clinicians should emphasize that numerous health benefits occur in the absence of weight loss and that maintenance of an active lifestyle will reduce the risk of future weight gain (Fogelholm and Kekkonen, 2000).

Table1: Expected Initial weight loss and possibly producing clinically significant Weight loss from different modalities of exercise training.

Modality	Weight loss	Clinically significant weight loss
Pedometer based step goal	Range: 0-1kg of weight loss	Unlikely
Aerobic exercise training only	Range: 0-2kg of weight loss	Possible, but only with extremely high exercise volumes
Resistance training only	None	Unlikely
Aerobic and Resistance training only	Range: 0-1kg of weight loss	Possible, but only with Extremely high exercise volumes of aerobic exercise training
Caloric restriction combined with aerobic exercise training	Range:-9kg to-13kg	Possible

Source: (Damon *et al.* 2014).

2.13. Exercise and Weight Maintenance

The ACSM position stand on PA intervention strategies to promote weight loss and weight regain emphasizes the distinction between the minimum levels of PA to maintain health (150 min/wk.) and higher levels of PA to prevent weight regain (200 min/wk.). Therefore, obese individuals who have successfully lost weight require a substantial amount of PA to maintain this weight loss. As indicated in the ACSM position stand (Donnelly *et al.*, 2009) several major limitations to research of PA on weight regain exist including the observational and the retrospective nature of the existing literature from randomized trials.

However, several studies in this area deserve mentioning. Using data from a PA weight loss study (Jakicic *et al.* 1999). observed a dose response between the amount of self-reported PA per week and long-term success with weight loss at 18 months of intervention (composed of caloric restriction and ET). Adults who exercised greater than 200 min/wk. (-13.1 kg) lost more weight compared to those who exercised between 150 and 199 min/wk. (-8.5 kg), and those that exercised less than 150 min/wk. (-3.5 kg). Different study by (Jakicic *et al.*, 2003)

observed similar findings in post hoc analyses of a weight loss intervention composed of both caloric restriction and exercise training in women. After 12 months of intervention, women with greater than 200 min/wk. (13.6%) had maintained significantly greater percentage of weight loss compared to those who had exercised at 150–199 min/wk. (9.5%), and less than 150 min/wk. (4.7%). Lastly, (Andersen et al. 1999) evaluated the effect of low-fat diet (1200 kcal/day) in combination with either structured aerobic ET or lifestyle activity (patients were advised to increase their PA to recommended levels) and both groups lost approximately 8 kg of weight following 16 weeks of intervention.

Weight maintenance was monitored for 1 year after the intervention and those who were the most active lost additional weight (1.9 kg) whereas the group that was the least active regained a substantial amount of weight (4.9 kg). These data suggest that PA has an important role in the amount of weight regain following successful weight loss. Clinicians should therefore advocate that their patients attempting to reduce recidivism after weight loss engages in PA levels above 200 min/wk. (Donnell *et al.*, 2009).

2.14. Sedentary Lifestyle and its Effects

2.14.1. Sedentary Lifestyle

A sedentary lifestyle is a type of lifestyle with no or irregular physical activity. It is commonly found in both the developed and developing world. Generally, people who have a sedentary lifestyle spend time at the computer, watching TV, reading or sitting. They spend almost no time getting in any kind of physical activity and involve low levels of energy expenditure (Market *et al.*, 2010).

2.14.2. Effects of Sedentary Life Style

A sedentary lifestyle and extended dose of sedentary behavior can result in dramatically increased metabolic risks and contribute too many preventable causes of death. (Hamburg *et al* 2007). One of the main effects of sedentary lifestyles is becoming at high risk to diseases and illnesses, like ischemic heart attack and chronic hypertension. This is usually due to lack of exercise which will lead to increase in cholesterol level and blood pressure. Sedentary lifestyles can also cause deep vein thrombosis, which is a condition that involves a blood clot formed in one of your veins, usually in the legs. This may not be very disturbing at first, but it

can be fatal once the clot dislodges and becomes an obstruction in the blood circulation of your heart and lungs. Another major effect of sedentary lifestyles is obesity.

Obesity can also lead to cardiovascular disorders, because of increased cholesterol level, and other neurological condition, such as stroke. This can also lead to even more medical condition such as fractures and muscle injury. The increase in weight of an obese person who seldom or does not exercise can make his bones prone to bone breakage. One fatal effect of this is when a broken part of the bone suddenly enters the bloodstream and becomes an obstruction to the blood circulation of the heart, lungs or brain. Low self-esteem can also happen if the person is disturbed by his increase in weight and body fat. Low energy level can also be one of the effects of sedentary lifestyles. Experts say that regular aerobic exercise can increase a person's energy level and can help the body and mind become.

2.15. Contributing Effects of Aerobic Exercise Intensity or Resistance Training on Weight loss ?

Vigorous intensity aerobic ET has been shown to have enhanced health benefits for important risk factors, including visceral fat, (Irving *et al.*,2008) measures of glucose/insulin metabolism, and CRF, (O'Donovan *et al.*,2005) compared to moderate intensity aerobic ET. In terms of weight change, when different intensities of ET are matched for caloric expenditure or ET dose, both vigorous and moderate intensity aerobic ET result in similar amounts of weight loss. O'Donovan .in an RCT of 64 obese men observed similar changes in weight following 6 months of moderate intensity (-1.1 kg) and high intensity (-0.5 kg) aerobic ET. In the STRRIDE study,(Kraus *et al.*,2002)observed similar weight loss in overweight/obese adults participating in 8 months of moderate (-0.6 kg) and high intensity (-0.2 kg)aerobic ET at the same exercise dose (14 kcal per kg per week).Vigorous intensity aerobic ET can contribute to greater weight loss if matched for session time compared to moderate intensity aerobic ET because the total energy expenditure is greater. Thus, if vigorous aerobic ET can be sustained and is enjoyable for the patient, the exercise program may induce additional health benefits and potentially increase the energy expenditure seen with exercise training (if replacing the exercise time of moderate intensity training).Resistance ET and health benefits including increasing/maintaining muscular strength with aging (prevention of sarcopenia) and

preserving bone mineral density. (Haskell *et al.*, 2007) Although, resistance ET alone contributes to the reduction of body fat, the effect on overall weight loss is minimal. (Donnelly *et al.* 2009) In the Health Benefits of Aerobic and Resistance Training (HART-D) study, (Church *et al.* 2010) observed no significant change (-0.3 kg) in weight in their assistance ET group (n = 73) compared to a control group after the 9 month intervention. In the DARE trial, no significant difference was observed between the resistance ET group and the control group (0.3 kg). (n = 86) observed no significant change in weight (0.07 kg) following 8 months of in the STRIDDE AT/RT study. (Bateman *et al.* 2011) Overall, little evidence exists that resistance training alone promotes weight loss.

CHAPTER THREE

3. RESEARCH METHODS


This chapter includes description of experimental site study period definition of variable, experimental design, source of data, study population, sampling size and sampling techniques, inclusion and exclusion criteria, data collection instrument, method and procedures of data collection, body weight, waist circumference description, aerobics training protocol, data quality control, methods of data analysis and ethical consideration.

3.1. Study Area

The practical experiment is conducted at Shashemene City which is located in the West Arsi 250km far from Addis Ababa. The experimental site is located at 38⁰ 36'E longitude and 7⁰12'N latitude and 1990m altitude above sea level In the Wester Ethiopia. Based on the 2007 Census conducted by the Central Statistical Agency of Ethiopia (CSA), Shashemene City has a total population of 100,454 whom 50,654were men and 49,800were women.

Shashemene 25 kilometers north of Hawassa City and 25km away from Negale Arsi Town 27 km from Kofale town 15km from Wondo Genet and 25km from shalla town. The climate of the State is one of the most pleasant in the country. Temperature is even between 12°C-7°C throughout the year. The coolest season (10°C) which is between June-October, coincides with heavy rains accompanied by storms and strong electrical discharges. The average annual intensity of precipitation is about 24,700 mm. The mean amount of rainfall over three years as registered 30years ago was 1.509 mm. This region is the only one in Ethiopia where the majority of its population lives in urban area: 272,084 or13.85% of the population are urban inhabitants. With an estimated area of 2,448square kilometers, this region has an estimated density of 24,700 people per square kilometer. For the entire region 387,143 households were counted, which results in an average for the Region of5.01 persons to a household, with urban households having on average 4.3 and rural households 5.2 people, CSAE. (2004).

Figure 2. Map of The Study Site



Source: Ethio-GIS 2015

3.2. The study design

In this research study the Pre-test and post-test randomized group pre-experimental design was used. 40 subjects were classified equally into two groups randomly. The experimental group was involved in an aerobics training program for 12 weeks. The control group was not participated on exercise because of comparing the effects of exercise with the treatment group. Pre-test and post-test were administered for both groups. A Pre-post-test of body weight and waist circumference test was taken from both groups based on the standard testing protocol. Aerobics exercise training was delivered for 60 minutes and 3days in a week for 12 weeks of training period (Monday, Wednesday, Friday)with moderate intensity (55-69% Vo_2 max).

3.3. Population of the study

Study population of women in Shashemene city was 49,800. There were fitness centers in the city that has been giving service for overweight people to address their goals. From the mentioned fitness center of its facilities Shashemene Secondary school. First, notice was posted for sedentary women in the city. Next, registration process was administered 150 women were registered. Then the investigator was made subjects to fill physical activity readiness questionnaires to identify and reject subjects with health problems. Finally forty subjects were selected from the total subjects and they were assigned in to two groups equally using systematic random sampling technique.

3.4. Sample Size and Sampling Techniques

Multi stage sampling techniques were used to specify study subjects. 150 volunteer overweight women filled the medical history questionnaire. The questionnaire was prepared with the aim of identifying whether they were free from cancer, heart disease, stroke and kidney problems. Additionally, injury statuses were used as one of the selection criteria. Finally, forty (40) subjects were selected for study and they were assigned into two groups equally using random sampling techniques. Those are the experimental group and control group. This means twenty(20) for the exercise group and twenty(20) for the non-exercise group which was used to compare with the experimental group to measure the effect of aerobics exercise among the group. Selected criteria was the following method:

Stage1. 150 voluntary overweight women registered to participate to this study.

Stage2. From 150 only 110 were came to fill the medical history questionnaire.

Stage3. The questionnaire was prepared with aim of identifying whether they were free from

Chronic disease: Like cancer, heart disease, and hypertension and kidney problem.

Stage 4. Injury statuses were used as one of selection criteria

Stage 5. From 70 voluntary overweight women were rejected due to factors listed in stage 3.

Stage 6. Finally 40 subjects were selected purposively for study and they were assigned equally in to experimental group and control group randomly by using lottery method.

3.5 Inclusion and Exclusion Criteria

Forty women those age ranges between 25 and 35 were used as a subject for the study. The participants should be free from any impairments, disability and chronic disease which members of Shashemene community and volunteer to respond to the designed study were included. Participants who were not eligible for the inclusive criteria were excluded. Women with different factors that affect their performance were excluded. Physical activity readiness questionnaire were used to include and exclude subjects.

3.6. Source of the data

Primary data were used in this research study. The primary data were obtained from the experimental variables according to the designed parameters

3.7 Data Collection Instruments

The whole data collection were performed more of quantitatively. Machines of the gymnasium like treadmill, cycles, step up box; digital weighting machine and meter were used during research investigation. The use of these principal data collection instrument was intended to explore a range of quantitative information. The researcher was used writing pad is also used to record the data

3.8 Method and Procedures of Data Collection

The experimental test was strictly administered and standardized in terms of administration, organization and implementation conditions. Up on starting the training programs pre-test was made. Then after the intervention the end of three months post-test were done.

3.8.1 Body Weight

Body weight test was taken using the digital weighting machine based on the administration protocol to know the initial base line and to observe the changes that was occurred in intervention. The height of subjects was taken using measuring tape.

Finally the body mass index was calculated as the following formula.

BMI= (W/H^2) where W= weight in kilogram, and H= height in meter²

Adult Body Mass Index or BMI

Body Mass Index (BMI) is a person's weight in kilograms divided by the square of height in meters. A high BMI can be an indicator of high body fatness and having a low BMI can be an indicator of having too low body fatness. BMI can be used as a screening tool but is not diagnostic of the body fatness or health of an individual.

To calculate your BMI, see the BMI Calculator. Or determine your BMI by finding your height and weight in this BMI Index Chart 1.

If your BMI is less than 18.5, it falls within the underweight range.

If your BMI is 18.5 to 24.9, it falls within the normal or Healthy Weight range.

If your BMI is 25.0 to 29.9, it falls within the overweight range.

If your BMI is 30.0 or higher, it falls within the obese range.

3.8.2 Waist Circumference Description

Waist Circumference is a measurement taken around the abdomen at the level of the umbilicus (belly button). Health experts use waist circumference to screen patients for possible weight-related health problems. If your waist measurement is greater than the numbers indicated, your risk for weight-related health problems is higher than normal.

The accurate measurement of waist circumference is achieved by using the following techniques:-

Locate the top of the hip (iliac crest) and take the measurement just above this bony landmark, just where one finger can fit between the iliac crest and the lowest rib. Ensure that the tip measure is positioned horizontally, parallel to the floor. Measuring at a level just above the iliac crest, and positioned the tape of irrespective of whether umbilicus is above or below the tape, provides the correct waist circumference measurement and should correspond to the maximum abdominal diameter. Ensure that the person is standing erect and has relaxed the abdominal muscle measurement is taken at the end of normal expiration.

Aim to have a snug but not tight a fit of the tape measure around your waist, do not make compressions in the skin with the tape measure. Accuracy can be approved by using a specially designed abdominal circumference tape measure. A constant-tension spring-loaded tape device reduces errors for over-enthusiastic tightening during measurement and improves accuracy and consistency of serial measurements.

The instrument we use to measure waist circumference is only meter.

Norem for waist circumference in adults		
Waist circumferences		
Risk category	Females	Males
Very low	<27.5 in (< 70cm)	<31.5 in (< 80cm)
Low	27.5-35.0 in(70-89cm)	31.5- 39.0 in (80-99cm)
High	35.5- 43.0 in (90-109cm)	39.5- 47.0 in (100-120cm)
Very high	>43.5 in (> 110 cm)	>47.0 in (>120 cm)

Source: Bray (2004).

3. 9. Aerobics Training Protocol

The selected subjects were participated on moderate intensity 55-69% HRmax aerobics training program. The duration of the study was for three consecutive months from October, 2021-December, 2021. For the first two consecutive weeks the experimental groups of participants were involved in aerobics exercise with low intensity (~40-50% HRmax) as a physical preparation or readiness, but the rest ten weeks experimental group were engaged with moderate intensity training program. The training program consisted of different physical exercises such as warming up, light stretching, brisk walking, running, cycling, aerobics dance, sit ups, push up, steeps and others. The frequencies of the days were three times in a week (Monday, Wednesday and Friday) from 4pm-5pm in moderate intensity exercise training program.

3.10. Data Quality Control

The quality of the data was assured by giving pre training for the subjects during data collection by the investigator. The researcher will check the collected data for completeness and on spot corrective measures on the field was taken accordingly.

3.11 Method of Data Analysis

Descriptive statistics will calculate the mean and standard deviations, in which the data was analyzed by SPSS version 20 statistical software package. It was analyzed through paired T-test was used to compare the pre training and post training dat. Level of significance less than 0.05 was considered.

3.12 Ethical Issues and Code of Conduct

The study was deal with the ethical issues, it will heed for the privacy of research participants and make guarantees and confidentiality in risk of harm as a result of their participation. Therefore, the study was conducted according to Jimma University rules, regulations, policies and codes of ethics relating to research ethics. Ethical standards require that researcher should not impose participants in a situation where they might be at risk of physical or psychological harm as a result of their participation. This research study was approved by an Ethics Review Committee of the Jimma University and Jimma Campus to make sure it do not result any risk or harm to the participants of this study.

CHAPTER FOUR

4. RESULTS AND DISCUSSION

4.1 DISCUSSION

This chapter deals with the analysis of pre and post test data collected from randomly selected experimental (n=20) and control (n=20) groups under the study. The purpose of this study was to investigate the effect of 12 week aerobic exercise program on the weight loss of sedentary overweight women in Shashemene city. Selected three variables (Bodyweight, Body mass index and waist circumference) and their component had been measured. Aerobic exercise was given for 12 weeks and attendance was taken for experimental throughout the exercise. Pre-test and post-tests were taken from both experimental and control groups before and after 12 weeks of aerobic exercise intervention and the scores were recorded. The collected data were analyzed using paired t-test to analyze pre-test and post-test results of experimental and control groups.

4.2 Results of the Study

4.2.1 Characteristics of study participant variables

As indicated in table 2 the mean values of participants' Age were before training m= 30.15 and after training m=30.15 for control group. Before training m=30.50 and after training for experimental group respectively. The mean value of body weight before training was m=72.15 and after training was m=73.25 for control group. The mean value of body weight before training was m= 71.20 and after training m= 67.15 for experimental group. The deference mean of body weight was increased by -1.1 for control group and reduced 4.05 for experimental group throughout the study period. The mean value of waist circumference before training was m=90.80 and after training was m= 91.75 fore control group and for experimental group before training m=91.45 and after training was 82.90. The mean of WC was increased by 0.95 and 8.55 was reduced experimental group throughout the study period. The mean value of body mass index before training was m=26.51 and after training was 26.92 for control group. Fore experimental group before training m=26.233 and after training

m=24.75. The mean of body mass index was increased by m=-0.04 fore control and fore experimental decrease by 1.48. This result was consistent with the finding of Willis and his friends. The result of their investigation showed that Aerobic exercise was efficient method of exercise for losing body weight and body fat (Willis *et al.*, 2012).

Table 2 The mean value of body weight, height, age and body mass index of both Controls group and experimental group.

Control group				Experimental group	
Variable		Pre	Pot	Pre	Pot
Ag	Mean	30.15+2.06	30.15+2.06	30.50+2.965	30.50+2.965
HT		1.65+0.023	1.65+0.005	1.65+0.029	1.65+0.006
BW		72.15+1.871	73.25+2.712	71.20+1.704	67.15+1.814
BMI		26.519+0.772	26.921+0.533	26.233+0.533	24.751+0.653

Values are mean ± SD, PT= pre training test which was taken before training

*POT= posttest which is taken after 12th week of aerobics training, WT =Weight, HT =height, AGE= age
BMI=body mass index*

As indicate in the below table 3 the pretest score of CG (M=90.80, MD=0.95, SD=0.76, T= -5.596, P=0.000 and the value of EG (M=91.45, MD= -8.55, SD= 2.39, T=15.98, P=0.000 Where As Post test Score of CG (M= 91.75, MD=0.950, SD=0.76, T= -5.56, P=0.000 and EG (M=82.90, MD= -8.55, SD=2.394, T=15.97, P=0.000. During pretest the mean value of CG and EG was nearly the same. In contrast after 12 week aerobic training the mean value of CG and EG was different. From this data can easily understand that the mean value of EG showed a statistical significant improvement on waist circumference (P< 0.05) rather than CG.

Table 3. Mean values of Body weight (kg), Waist circumference (WC) of (CG) and (EG); at different occasions of training program.

Var	Control group		Experimental group		P-Value
	PT	Pot	PT	Pot	
BWT	72.15+1.87	73.25+2.71	71.20+1.70	67.15+1.81	0.00
WC	90.80+1.641	91.75+1.80	91.45+1.67	82.90+2.46	0.00

Values are mean + SD, CG= control group, EG= experimental group, PT= pre training test which is taken before training and POT= post training test measured at the 12th week of after training and BWT= body weight and WC= waist circumference.

Paired sample t-test results of each parameter

As indicate in the belo table 3 the pretest score of CG (M=72.15, MD= -1.100, SD=1.209 T=-4.067, P=0.001 and the value of EG (M=71.20, MD=4.050, SD= 0.759, T=23.85, P=0.000). Where as the post test score of CG (M=73.25, MD=-1.100, SD=1.209,T=-4.067, P=0.001 and the value of EG (M=67.15, MD=4.050, SD=0.759, T=23.858, P=0.000). During pretest the mean value of CG and EG was nearly the same. In contrast after 12 week aerobic training the mean value of CG and EG was different. From this data can easily understand that the mean value of EG showed a statistical significant improvement on body weight (P< 0.05) rather than CG.

Table 4. Paired t test results of body weight

	Tests	Mean	MD	SD	Lower Bound	Upper Bound	T value	P value
CG	Pt-pot	72.15-73.25	-1.100	1.209	-1.671	-533	-4.067	0.001
EG	Pt-pot	71.20-67.15	4.050	0.759	3.694	4.405	23.858	0.000

MD=mean difference PT = pre training test which was taken before resistance training, WC= Waist Circumference, POT= posttest which is taken after 12th week of aerobic exercise training. WT =Weight, BMI=body mass index

As indicated in the below table5 the pretest score of CG (M=26.51,MD=-0.402,SD=0.45,T=4.004, P=0.001 and the value of EG (M=26.23, MD=-1.48, SD=0.061,T=23.96,P=0.000 where as the post test score of CG (M=91.75, MD=-0.402, SD=0.45,T=-4.004 P=0.001and EG (M=24.75, MD=-1.482, SD=0.061,T=23.96,P=0.000. During pretest the mean value of CG and EG was nearly the same. In contrast after 12 week aerobic training the mean value of CG and EG was different. From this data can easily understand that the mean value of EG showed a statistical significant improvement on body mass index (P< 0.05) rather than CG.

Table 5. Paired t test results of BMI

	Tests	Mean	MD	SD	Lower Bound	Upper bound	T value	P value
CG	Pt-pot	26.51-26.92	-0.402	0.45	-0.612	-0.191	-4.004	0.001
EG	Pt-pot	26.23-24.75	-1.482	0.061	1.353	1.612	23.960	0.000

MD=mean difference PT = pre training test which was taken before resistance training, WC= Waist Circumference, POT= posttest which is taken after 12th week of aerobic exercise training. WT =Weight, BMI=body mass index

As indicate in the below table 6 the pretest score of CG (M=90.80, MD=0.95, SD=0.76,

T= -5.596, P=0.000 and the value of EG (M=91.45, MD= -8.55, SD= 2.39,T=15.98, P=0.000

Where As Post test Score of CG (M= 91.75, MD=0.950, SD=0.76,T=-5.56, P=0.000 and EG

(M=82.90, MD= -8.55, SD=2.394, T=15.97, P=0.000. During pretest the mean value of CG

and EG was nearly the same. In contrast after 12 week aerobic training the mean value of CG

and EG was different. From this data can easily understand that the mean value of EG showed

a statistical significant improvement on waist circumference (P< 0.05) rather than CG.

Table 6. Paired t test results of Waist Circumference

	Tests	Mean	MD	SD	Lower Bound	Upper Bound	T value	P value
CG	Pt-pot	90.80-91.75	.950	0.759	-1.305	-0.594	-5.596	0.000
EG	Pt-Pot	91.45+82.90	-8.55	2.394	7.429	9.670	15.968	0.000

MD=mean difference PT = pre training test which was taken before resistance training, WC= Waist Circumference, POT= posttest which is taken after 12th week of aerobic exercise training. WT =Weight, BMI=body mass inde

A positive mean difference was observed for experimental group. The rationale behind the decrease of body weight for experimental group was due to aerobics training they took in the aerobic exercise. The mean value of body weight before training was 72.15 and 73.25after training for control group. The mean value of bodyweight before training was 71.20 and 67.15 after training for experimental group. The mean of body weight was increased by 0.76 for control group and 2.39 was reduced in experimental group throughout the study period. The mean value of waist circumference before training was 90.80 and 91.75 after training for control group. The mean value of waist circumference before training was 91.45 and 82.90 after training fore experimental group. The mean of waist circumference was increased by – 0.95 fore control group and 8.55 was reduced in experimental group throughout the the study

period. In the experimental group weight of body of the participants were reduced significantly due to aerobics training. This result was consistent with the finding of Willis and his friends. The result of their investigation showed that Aerobic exercise was efficient method of exercise for losing body weight and body fat (Willis *et al.*, 2012). This result was also agreed with finding of Arslan. He pointed out eight weeks step aerobic dance exercise significantly decreased body composition parameter of middle aged sedentary obese women (Arslan, 2011) Aerobic exercise including walking, running, and swimming has been proven to be an effective way to lose weight. In waist circumference test, the mean value of control group was increased but in experimental group waist circumference was decreased markedly. The mean difference showed that there was improvement in the EG. The EG was decreased 8.55 but -0.95 was increased for the CG.

The aerobic exercise significantly reduced the body mass index and body weight improved cardiovascular endurance, abdominal endurance/strength and flexibility of individuals. Similarly, in a study conducted by (Hopkins *et al.*, 1990), proved significant improvement in all functional fitness components of their participants including cardio respiratory endurance, strength/endurance, body agility, flexibility, body fat and balance was reported. This study was also corroborated by (Petrofsky *et al.*, 2008) who did their study on the effect of aerobic dance and diet program on cardiovascular fitness, body composition, and weight loss in women. They reported a significant decrease in body weight, reduced waist girth and an improved cardiovascular function and general fitness.

The mean value of body mass index (BMI) was increased from one test to another for control group. And more significant change distinguished decreased for the EG. The EG exhibited 5.05% decreased in BMI. This result was in agreement with the findings of (Leijssen *et al.*, 2002) noted that aerobics exercise can be an important component of weight lose intervention and therefore, commonly included as part of comprehensive weight lose management program. Additionally, BMI was 0.95% increased recorded in the CG.

Table 7: The mean difference value and significance level of each test results in both EG and CG

Dependent variable	Para I	Para II	MD(II-I)		P Value	
			EG	CG	EG	CG
Body weight (kg)	PT	POT	4.050	1.100	.000	.000
BMI (kg/m ²)	PT	POT	-1.300	.250	.000	0.16
Waist circumference(WC)	PT	POT	-8.550	.950	.000	.000

MD=mean difference PT = pre training test which was taken before resistance training, WC= Waist Circumference, POT= posttest which is taken after 12th week of aerobic exercise training. WT =Weight, BMI=body mass index

CHAPTER FIVE

5. SUMMARY, CONCLUSION, AND RECOMMENDATION

5.1. Summary of the major findings

The purpose of this study is to find out the effect of 12weeks aerobic exercise program on reducing body weight of sedentary over weight women in the community of Shashemene City. To achieve the purpose of this study 40 sedentary female and grouped into experimental and control group equally to compare the effects of the given exercise. Both groups were selected from which age ranges 25-35 years old. The exercise program was designed for 12 weeks, three(3) times per week with 60-minute duration and moderate intensity. Each session was divided again in to warming up, main part (aerobic exercise stretching) and cooling down phase. The data collected from the study was analyzed using SPSS version 20 software. The paired sample t-test was used for this study. The mean values of participants' age were 30.15 and 30.50 for control group and experimental group respectively. The mean values of participants' height were 1.65 and 1.65 for control group and experimental group respectively.

The mean value of body weight before training was 72.15 and 73.25 after training for control group. The mean value of body weight before training was 71.20 and 67.15 after training for experimental group. The mean of body weight was increased by 1.52% for control group and 5.688% was reduced in experimental group throughout the study period. In the experimental group weight of body of the participants were reduced significantly due to aerobics training. In waist circumference before training was 90.80 and after training was 91.75 fore control and fore experimental before training was 91.45 and after training was 82.90. The mean value of control group was increased but in experimental group waist circumference was decreased markedly. The mean difference showed that there was improvement in the EG. The EG was decreased 9.34% but 1.04% was increased for the CG. The mean value of body mass index was 26.52 before training and after training was 26.92 fore control group. The mean value of body mass index before training was 26.23 and after training was 24.75 fore experimental group. Body mass index increase by -0.042 fore control and 1.48 decreased by 1.48 fore experimental group. The test measurements used to identify the improvements BW, BMI, and

WC. Based on the analysis made, at the end of the program it was observed that body weight, waist circumference and BMI reduced significantly for exercise group (Experimental). But control group was increased in the measured parameters.

The reduction of the experimental group in the measured variables was due to the given exercise training program and highly significant change was observed in improvement of body weight, waist circumference, and body mass index for experimental group.

5.2. Conclusions

Depending on the major findings of this study, the following points were stated as a conclusion:

- All parameters clearly showed that the better test results were recorded in post-training than pre training. The mean of body weight was reduced by 4.05 in experimental group throughout the study period. This indicates that moderate intensity aerobics exercise training program were effective for the reduction of body weight.
- Continuing participating in moderate intensity aerobics exercise program had appositve effect on body mass index. Body mass index decreased by 1.48 resoult.
- The mean difference showed that there was improvement in the EG. The EG was decreased by 8.55, aerobic exercise training program had a positive effect on waist circumference.
- In general, this finding clearly noted that moderate-intensity aerobics exercise training program has a significant effect on the reduction waist circumference body weight and body mass index that resulted following the accumulation of excessive fat in the body. Aerobic exercises are interlinked.

5.3. Recommendations

The findings of the study showed that a 12 week of aerobic exercise program can have a significant effect on body weight reduction and waist circumference improvement for overweight people. Based on the findings, the following recommendations are made:

- It is highly expected from professionals of sport science, and related fields to guide and educate on the importance and value of aerobic exercise on reduction of body weight for health benefits.
- In view of the benefits of aerobic exercises educational authorities may consider inclusion of this exercise as a part of the aerobic exercise program for all over weight woman
- Since aerobic exercise require no equipment or minimal equipment. Sedentary female's community may be encouraged to undergo this type of exercise training regularly for physical fitness improvement.
- It is good, if government and private organizations introduce one hour aerobic exercise program three times weekly for their staff which may help in improving work force and efficiency among workers
- It is advisable that the government should create a forum for seminar and workshop on various aerobic exercise programs and their importance and benefit to health.
- This all above can help to reduce the possibility of obesity, overweight and improves cardiovascular endurance, muscular strength/endurance and flexibility.

REFERENCES

- Andersen, R.E., Wadden, T. A. and Bartlett, S.J.1999. Effects of lifestyle activity versus structured aerobic exercise in obese women: a randomized trial. *JAMA*;281:335-340.
- Arslan, J., 2011. Conducted on the study on the effects of an eight-week step-aerobic dance Exercise program on body composition parameters in middle-aged sedentary obese women.*International Sport Med Journal*, 12 (4):160-168
- Ays, E., Sarsan, F. Sun, A. Merih, O.andZgen, O. T.2006.The effects of aerobic and resistance exercises in obese women.*Clinical Rehabilitation*.
- Bateman, L. A.,Slentz, C. A.and Willis L. H. 2011. Comparison of aerobic versus resistance exercise training effects on metabolic syndrome (from the Studies of a Targeted Risk Reduction Intervention Through Defined Exercise - STRRIDE-AT/RT). *Am J Cardiol.*;108:838-844.
- Behnke, A. R., J. H. Wilmore,1974. Evaluation and Regulation of Body Build and Composition. Englewood Cliffs, NJ: PrenticeHall.
- Brodie, D. A.,1988. Techniques of measurement of body composition: Part I. *Sports Medicine*. 5:11–40.
- Brown, W.C., I.A. Dubuque, 1986. Physiology of exercise for Physical Education and athletic, 4thedition. P. 255-260
- Castaneda, C., Layne, J.E. Munoz,O. L. Gordon,P .L.Walsmith, J. Foldvari, M.Roubenoff, R. Tucker, K.L. and Nelson, M.E.2002. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care*, 25: 2335–2341.
- Church, T.S., Blair, S. N. andCocreham, S. 2010. Effects of aerobic and resistance training on hemoglobin a1c levels in patients with type 2 diabetes. *JAMA.*;304:2253-2262.

- Church, T.S., Earnest, C.P. and Thompson, A.M. 2010. Exercise without weight loss does not reduce C-reactive protein: The INFLAME study. *Med Sci Sports Exerc.*;42:708-716.
- Church, T.S., Earnest, C.P. and Skinner, J.S. 2007. Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure. *JAMA.*;297:2081-2091.
- Church, T. S., Martin, C. K. and Thompson, A.M. 2009. Changes in weight, waist circumference and compensatory responses with different doses of exercise among sedentary, overweight post-menopausal women.
- Cooper, K.H., 1968. A means of assessing maximal oxygen uptake. *Journal of the American Medical Association*, 203:201-204
- Corbett, E., 2009. Effects of oral L- Arginine supplementation on platelet count and maximal oxygen consumption in healthy males. The graduate faculty of the University of Akron. pp.11
- Damon, L., Swifta, B. Neil, M. Johannsenc, F. Carl, J. Laviec, F. Conrad, P. Earnestd, Timothy, S. and Church .2014. The Role of Exercise and Physical Activity in Weight Loss And Maintenance. *Progress In Cardiovascular Diseases* 56, 441 – 447.
- DiPietro, L., Dziura, J. and Yeckel, C.W. 2006. Exercise and improved insulin sensitivity in older women: Evidence of the enduring benefits of higher intensity training. *Journal of Applied Physiology*, 100:142–9.
- Donnelly, J., E. Blair, S. N. and Jakicic, J. M. 2009. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc.*;41:459-471.
- Donnelly, J., E. Blair, S.N. and Jakicic, J.M. 2009. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc.*;41:459-471.

- Evrin, C., S. Ahmet, V. Husamettin, and A. Kamile, 2010. The effect of 8-week step-aerobic exercise on the body composition and hematologic parameters in the obese and overweight females. *The New Sports of Medicine*, 15 (1):14
- Sharon, A., Plowman, Denise, and L. Smith, 2011. *Exercise physiology for health, fitness, and performance*. —3rd ed. Philadelphia, PA 19106.
- Fitwatch, and Shemels, M. 2010. workoutplanner. www.fitwach.com/planner.
- Fogelholm, M., Kukkonen-Harjula, K. 2000. Does physical activity prevent weight gain—a systematic review. *Obes Rev.*;1:95-111.
- Fogelholm, M., and Kukkonen, H. K. 2000. Does physical activity prevent weight gain? *Medical Journal*, 1:95–111.
- George, A., Kristi, S. Kelley, Susan, R. and William, H. 2012. Combined effects of aerobic exercise and diet on lipids and lipoproteins in overweight and obese adults: a meta-analysis.
- Gian, P.E. Maria, C.G. Marco, M. Luigi, D. Silvia, M. Lorenzo, M.D. Felice, S. and Laura, G. 2015. Effects of Aerobic Exercise Based upon Heart Rate at Aerobic Threshold in Obese Elderly Subjects with Type 2 Diabetes. *International Journal of Endocrinology*.
- Golding, 1986. *The Y's way to physical fitness*. 3rd edition. Pp. 85-97
- Gutin, B. A., 1980. A Model of physical fitness and dynamic health. *Journal Health, Physical Education, and Recreation*, 51:48.
- Hamburg, N.M., C.J. McMackin, A.L. Huang, S.M. Shenouda, M.E. and Widlansky, 2007. Physical inactivity rapidly induces insulin resistance and micro vascular dysfunction in healthy volunteers. *ArteriosclerThrombVasc.Biol.*, 27 (12):2650–2656.

- Haskell, W. L., Lee, I. M. and Pate, R. R. 2007. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc.*;39:1423-1434.
- Hopkins, D.K., B.M. Murrain and K.C. Rhodes, 1990. Effect of Low Impact Aerobic Dance on the Functional Fitness of Elderly Women. *American Journals of Gerontology*, 30 (2):189-192
- Irving, B.A., Davis, C.K. and Brock, D.W. 2008. Effect of exercise training intensity on abdominal visceral fat and body composition. *Med Sci Sports Exerc.*;40:1863-1872.
- Jakicic, J.M., Marcus, B. H. and Gallagher K. I. 2003. Effect of exercise duration and intensity on weight loss in overweight, sedentary women: a randomized trial. *JAMA.*;290:1323-1330.
- Jakicic, J. M., Winters, C. and Lang, W. 1999. Effects of intermittent exercise and use of home exercise equipment on adherence, weight loss, and fitness in overweight women: a randomized trial. *JAMA.*;282:1554-1560.
- Jakicic, J.M., Clark, K. Coleman, E. Donnelly, J. E. Foreyt, J. Melanson, E. Volek, J. and Volpe, S.L. 2001. Appropriate intervention strategies for weight loss and prevention of weight regain for adults. *Medical Science of Sports Exercise*, 33: 2145–2156.
- Joav, M., Asad, B. Eli, C. and Einat, K. 2013. Effects of Aerobic Exercise on Body Composition and Muscle Strength in Over-Weight to Obese Old Women with Intellectual Disability: A Pilot Study *Open Rehabilitation Journal*, 6, 43-48
- Johannsen, N. M., Swift, D.L. and Lavie, C. J. 2013. Categorical analysis of the impact of aerobic and resistance exercise training, alone and in combination, on cardiorespiratory fitness levels in patients with type 2 diabetes mellitus: Results from the HART-D study. *Diabetes Care.*; 36(10):3305-3312.

- Kaminsky, L., and G. Dwyer, 2006 .Resource Manual for Guidelines for Exercise Testing and Prescription (th edition).Body Composition. In L.Kaminsky. (ed.), A CSM's Philadelphia, PA: Lippincott Williams and Wilkins, 195–205.
- Katch, V.L., D.M. William, and I.K. Frank, 2011. Essential of exercise physiology 4th Edition pp: 425
- Kathleen, K., 2006. Understanding the Benefits of Aerobic Exercise .The New England Wellness web
- King, N.A., Hopkins, M. and Caudwell, P.2008. Individual variability following 12 weeks of supervised exercise: Identification and characterization of compensation for exercise-induced weight loss. *International Journal of Obesity*, 32:177–84.
- Kraus, W.E., Houmard, J. A. andDuscha B. D. 2002. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med.*;347:1483-1492.
- Kraus, W.E., Houmard, J. A.andDuscha, B. D. 2002. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med.*;347:1483-1492.
- Kraus, W.E.,Houmard, J. A.andDuscha, B.D.2002.Effects of the amount and intensity of exercise on plasma lipoproteins. *Medical Journal*, 347:1483–92.
- Leijssen, V.A., W.Saris, A.J. Wagenmarks, J.M. Senden, and M.A.Baak, 2002.Effect of exercise training at different intensities on fat metabolism of obese men.*Journal of Applied Physiology*,92:1300-1139.
- Mark, S.T., H.N. Genevieve, C.C. Rachel, S.J. Travis, and O. Neville, 2010 .Physiological and health implications of a sedentary lifestyle. *Appl. Physiol. Nutr. Metab.*, 35: 725–740).
- Mills, and E. Mae, 1994.The effects of low intensity aerobic exercise on muscle strength, flexibility and change of balance among sedentary elderly persons. *Europe PMC; Nursing Research*, 43(4):207-11

- Musa Lewis M., 2013. The effect of a 10-week tae-bo intervention program on cardio metabolic disease risk factors in overweight and obese females at the University of Zululand, South Africa.
- O'Donovan, G., Owen, A. and Bird, S. R. 2005. Changes in cardiorespiratory fitness and coronary heart disease risk factors following 24 wk of moderate- or high-intensity exercise of equal energy cost. *Journal of Applied Physiology*;98:1619-1625.
- Odonovan, G., Owen, A. and Bird, S.R. 2005. Changes in cardio respiratory fitness and coronary heart disease risk factors following 24 week of moderate- or high-intensity exercise of equal energy cost. *Journal of Applied Physiology*, 98:1619–25.
- Ogden., C.L., Carroll, M.D. Curtin, L.R. McDowell, M.A. Tabak, C. J. and Flegal, K.M. 2006. Prevalence of overweight and obesity in the United States. *Journal of American Medicine Association*, 295: 1549–1555.
- Ozcan, S. and M. A. Ozturk, 2011. The Effect of twelve week aerobic exercise program on health related physical fitness components and blood lipids in obese girls. *Academic Journals*, 5 (12):1441-1445
- Pernille, N., Mads, R. Thorkil, P. Karina, W. Michael, F. Ninna, B. Nielsen, J. W. Helge, and Bente, S. 2015. Independent effects of endurance training and weight loss on peak fat oxidation in moderately overweight men: *Journal of Applied Physiology*. 118: 803–810.
- Petrofsky, J.B., J.B. Collins, T.L. Moein, K. Yang, M. Bains, D. Gunda, S. Raju, C. Vanaras, D. Kim, Y. Beard, C. Broussard, C. Christiansen, J. Ellstrom, C. George, I. Holland and J. Brown, 2008. Effect of Aerobic Dance and Diet Program on Cardiovascular Fitness, Body Composition and Weight Loss in Women. *Journals of applied research*, 8 (3):179-188
- Probart, C.K., M. Notelovtz, D. Martin, F.Y. Khan and C. Fields, 1991. The Effect of moderate Aerobic Exercise on Physical Fitness among Women 70 years and Older. *European Journal of Applied Physiology*, 14:49-56.

- Pu, L. I., Djoussé, L. and Sesso, H.D. 2010. Physical activity and weight gain prevention. *Medical Journal*, 303:1173–9.
- Ratamess, N. A., Chiarello, C. Sacco, A. J. Hoffman, J.R. Faigenbaum, A. D. Ross, R. E. and Kang, J. 2012. The Effects of Rest Interval Length on Acute Bench Press Performance: The Influence of Gender and Muscle Strength. *Journal of Strength & Conditioning Research: Volume 26 - Issue 7 - p 1817–1826*
- Roche, A. F., 1987. Some aspects of the criterion methods for the measurement of body composition. *Human Biology*. 59(2):209– 220 .
- Ross, R., Dagnone, D. and Jones, P. J. H. 2000. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. *Ann Intern Med.*;133: 92-103.
- Ross, R., Janssen, I. and Dawson, J. 2004. Exercise-induced reduction in obesity and insulin resistance in women: A randomized controlled trial. *Obes Res.*;12:789-798.
- Saris, W., Blair, S. N. and Van, B. 2003. How much physical activity is enough to prevent unhealthy weight gain? *Obese, Rev.*, 4:101–14.
- Selvalakshmi. 2007. Effect of varied aerobic training program on obese women working in IT companies, A Paper Presented at the National Seminar on Multidimensional Futuristic Approach to excellence in physical education and Sports; Pp. 42.
- Selvan, V.S. 2008. “Effect of Aerobic Exercise on Selected Physiological Variables among College girls” A Paper Presented at National Seminar on Professional and Scientific Approaches in Physical Education and Sports Sciences.
- Sigal, R.J., Kenny, G.P. and Boulé, N. G. 2007. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes. *Ann Intern Med.*;147:357-369.

- Suleen, S. H., Satvinder, S. Dhaliwal, A. P. Hills, and Sebely, 2012. The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial.
- Swift, D.L., Johannsen, N. M. and Lavie, C. J. 2014. The role of exercise and physical activity in weight loss and maintenance. *Prog Cardiovasc Dis*;56:441–7.
- Thomas, D. M., Bouchard, C. and Church, T. 2012. Why do individuals not lose more weight from an exercise intervention at a defined dose? An energy balance analysis. *Obes Rev.*;13:835-847.
- Thomas, D.F., M.I Paul, and T.R. Walton, 2008. FIT & WELL 8th Edition
- Thomas, D. 2012. Why do Individuals Not Lose More Weight from an Exercise Intervention at a Defined Dose? An Energy Balance Analysis. *Obesity Reviews*, 13, 835-47.
- Thomas, D.M. Bouchard, C. and Church, T. 2012. Why do individuals not lose more weight from an exercise intervention at a defined dose? *Medical Journal*, 13:835–47
- Trapp, E. G., Chisholm, D.J. and Freund, J. 2008. The effects of high-intensity intermittent exercise training on fat loss and fasting insulin levels of young women. *Int J Obes (Lond)*;32:684–91
- Van, W. G., Dubin, J.A. and Blair, S.N. 2007. Adult obesity does not predict 6-year weight gain in men. *Medical Journal*, 15:1571–7.
- Walter, R., Thompson, N. F. Gordon, L.S and Pescatello. 2010. Guidelines for exercise testing and prescription. 8th ed. American College of sport medicine.
- Wells, K.F., and E.F. Dillon, 1952. The sit and reach. A tests of back and leg flexibility. *Research Quarterly*, 23.115-118.
- WHO, 2003 Global strategy on Diet, Physical activity and Health. <http://www.who.int/dietphysicalactivity/media/en/gsfspa>. Accessed on October 20 (2012)

- William, R.G. Abdou, T. Samuel, H. M. Peguy, B. Assomo, N. Regine, T. Messina, O. André, H. Laurent, S. Etoundi, N. and Said, A. 2016. Effect of 24-week repeated short-time walking based training program on physical fitness of black Cameroonian obese women. *Journal of Exercise Rehabilitation*, 12(2):90-98.
- Williams, M. A., Haskell, W.L. Ades, P.A. Amsterdam, E.A. Bittner, V. Franklin, B.A. Gulanick, M. Laing, S.T. and Stewart, K.J. 2007. Resistance exercise in individuals with and without cardiovascular disease. *Journal of American Heart Association*, 116: 572–584.
- Willis, L. H., C.A. Sletzer, L.A. Bateman, A.T. Shields, L.W. Piner, C.W. Bales, J. A. Houmard and W. E. Kraus, 2012. Effects of aerobic training on body mass and Fat mass in overweight or obese adults. *Journal of Applied Physiology*, 113 (12):1831

APPENDICES
JIMMA UNIVERSITY

SPORT ACADEMY

DEPARTMENT OF SPORT SCIENCE

Appendix I Health History and Physical Readiness Questionnaires

This questionnaire will be applied to obtain information on health status and physical readiness of the trainers participating for research study entitled “Effects of 12 Weeks of Aerobic Exercises Program on Weight Loss of Sedentary Over Weight Women in Shashemene City”.

The information will be kept strictly confidential.

General information

Sex F Age----- phone-----

Weight ----- kg Height-----m

For participants; please read the following questions carefully and indicate your correct response for each question by encircling on the choice letter given below. Common sense your best guide when you answer this questions.

1. Has your doctor ever said that you have a heart condition and that you should only do physical activity until you recommended doing so?

A. Yes B. No

2. Do you feel pain in your chest when you do physical activity?

A. Yes B. No

3. In the past months have you had chest pain when you were not doing physical activity?

Purpose of the study:

Purpose of this study is to examine the effect of post supplementation of whey protein on body weight and muscle strength of sport Female science students using 12 weeks regular resistance exercise.

Procedure and duration:

You are being asked to participate in this research study as described below. All this like research study carried out are governed by the regulations for research on human beings. These regulations require that the researcher should obtain a signed agreement (consent) from you to participate in this research project.

The researcher had explained to you in detail the purpose of the project, the procedures to be used, the potential benefits and the possible risks of participation in this study. You can ask the researcher any questions that you may have about the study, and expect to receive satisfactory answers regarding the same. A basic explanation of the project is summarized below. After discussion, if you agree to participate in the study, please sign this form in the presence of the researcher. You may discontinue at any time from the study if you choose to do so.

Risks and benefits:

The risks of this research study are small. While administering the tests and during training session you may experience localized muscle fatigue in your muscles. You might feel some muscle soreness and fatigue during and after the cessation of the exercise tests and training but we do not expect any unusual risks as a direct result of this study. If any unexpected physical injury occurs, appropriate first aid will be provided, but no financial compensations will be given.

Confidentiality:

The information obtained about you will be kept in confidence, although you are free to release it to your own physician. The information will be used only for scientific purposes without identifying you as an individual.

Rights:

Participation for this study is fully voluntary. You have the right to declare to participate or not in this study. If you decide to participate, you have the right to withdraw from the study at any time and this will not label you for any loss of benefits which you otherwise are entitled. You do not have to answer any question that you do not want to answer.

Contacts address:

If there is any questions or enquires any time about the study or the procedures, please contact:

Ukula Yohannis Wabe at (+251922135811) or

E-mail: bontu Yoha@gmail.com. If you have any questions on your rights as a research subject, you can call the Institutional Research Ethics Review Committee of Jimma University.....

Appendix III Paired sample t test results of each parameter

1. Paired t test results of body weight

	Tests	Mean	MD	SD	Se	Lower Bound	Upper Bound	T value
Cg	Pt-pot	72.15-73.25	1.100	1.194	.189	.718	1.482	5.827
Eg	Pt-pot	71.20-67.15	-4.050	.749	.118	-4.290	-3.810	-34.182

2. Paired t test results of BMI

	Tests	Mean	MD	SD	Se	Lower Bound	Upper bound	T value
Cg	Pt-pot	26.51-26.92	0.41	.447	.100	-.557	-.275	-5.96
Eg	Pt-pot	26.23-24.75	-1.48	1.03	0.73	.174	-830	3.100

3. Paired t test results of Waist Circumference

	Tests	Mean	MD	SD	SE	Lower Bound	Upper Bound	T value
Cg	Pt-pot	90.80-91.75	.950	.749	.118	.710	1.190	-6.54
Eg	Pt-Pot	91.45+82.90	-8.55	4.84		2.42	5.479	5.22

Appendix IV Description of Aerobic Exercises Plan

Aerobic Exercise

Aerobic exercise helps strength your heart and lungs. Factors that affect aerobic exercise include how often you perform aerobic activity, the amount of time you spend at each session and the intensity (or percentage) of your maximum heart rate. Perform aerobic exercise:

3–5 times per week

20–60 minutes each time

at 60–85 percent of your maximum heart rate, OR

At 50–60 percent of your maximum heart rate if you are just starting an exercise program or have a heart condition at first, 3 times per week, for 20 minutes, at 60 percent of your heart rate may be enough activity for you. As your conditioning improves, increase the number of days per week, and the time and intensity of your exercise.(Sharon et al.,2011)

Maximum Heart Rate

A person's heart rate is the measure of heart beats per minute. Your maximum heart rate is the heart rate you should reach when exercising at a maximal exertion level. To calculate your maximum heart rate, subtract your age from 220. For example, if you are 30 years old, your maximum heart rate is 190: $(220 - 30 = 190)$

Target Heart Rate

Target heart rate is 50–85 percent of your maximum heart rate. Exercising within your target heart rate zone allows you to maximize your cardio respiratory fitness level. For example, if your maximum heart rate is 190, then the lower-end of your target heart rate zone is 95 heartbeats per minute $(0.50 \times 190 = 95)$.

If you are just starting an exercise program, or if you have a heart condition, your physician may recommend you begin exercising at 50–60 percent of your maximum heart rate. If you have been active, and do not have a heart condition, exercising at 60–65 percent of your

maximum heart rate may be a good place to begin. Once the target heart rate you choose becomes too easy for you, incrementally increase your target heart rate to the upper-end (85 percent) of your target heart rate zone ($0.85 \times 190 = 162$ heartbeats per minute). Determining your target heart rate requires periodic measuring. The best way to measure your heart rate is to purchase a heart rate monitor. The price of a good heart rate monitor can range from \$30 – \$100, but is worth the investment. They can be found at most sporting goods store (Sharon et al.2011).

Exercise Intensity Level

There are three levels of exercise intensity: low intensity, medium intensity, and high intensity. The intensity level of exercise determines your stage of exercise, and the amount of calories you burn each time. Identifying the appropriate exercise intensity level will help you determine the type of exercises you should perform (Sharon et al.2011).

Low Intensity:

For: beginners; those with a heart condition; those weighing over 250 pounds; or those new to med fast to exercise at a low intensity level, your target heart rate should be 50–65 percent of your maximum heart rate. To calculate your maximum heart rate, subtract your age from 220. For example, if you are 30 years old, your maximum heart rate is 190 ($220 - 30 = 190$). If you choose to exercise at 50 percent of your maximum heart rate, your target heart rate would be 95 heartbeats per minute ($0.50 \times 190 = 95$). If you choose to exercise at 65 percent of your maximum heart rate, your target heart rate would be 124 heartbeats per minute ($0.65 \times 190 = 124$). Depending on your weight low-intensity exercise will burn an average of 2.5 calories per minute. Begin incorporating low-intensity exercises for at least 20 minutes a day, 3–4 days a week. Gradually, work your way up to the next exercise level (medium-intensity). There are many low-intensity exercise options, including: Gardening raking leaves, pulling weeds, light shoveling etc. House work vacuuming, sweeping/mopping, cleaning window Walking Painting Washing the car (Sharon et al.,2011).

Medium Intensity:

For individuals comfortable with beginning a more active exercise program Exercising at a medium-intensity level requires the use of large muscle groups (such as the back, chest, legs, and buttocks). At a medium-intensity level, exercisers work at 65–75 percent of their maximum heart rate. To calculate your maximum heart rate, subtract your age from 220. For example, if you are 30 years old, your maximum heart rate is 190 ($220 - 30 = 190$). At a medium-intensity level, you should exercise anywhere between 124–143 beats per minute ($0.65 \times 190 = 124$; $0.75 \times 190 = 143$).

Medium-intensity exercise burns an average of 7.5 calories per minute depending on your weight. There are a wide variety of medium-intensity exercises you can do on a regular basis. These exercises are not only beneficial for losing weight – they also help prevent heart disease, lower blood pressure, reduce the risk of developing diabetes, and improve psychological well-being. Once you have become comfortable working out at a low-intensity level, incorporate medium-intensity exercises at least 3–4 days per week. Begin with a comfortable time frame of 15–20 minutes per day, and gradually work your way up to at least 45 minutes a day, 4–5 days per week. Examples of medium-intensity exercises include:

Brisk walking, Bicycling, Sports tennis, basketball, swimming, Dancing salsa, swing, Hiking etc, (Sharon et al, 2011).

High Intensity:

For regular exercisers (currently in Maintenance) looking to step things up High-intensity exercise is not for beginners or anyone just starting the med fast program. These exercises are recommended for individuals who exercise daily, and have either hit a plateau or are looking for something more intense to incorporate into their exercise routine. At a high intensity exercise level, you are exercising at 75–85 percent of your maximum heart rate. To calculate your maximum heart rate, subtract your age from 220. For example, if you are 30 years old, your maximum heart rate is 190 ($220 - 30 = 190$). At a high-intensity level, you should exercise anywhere between 143–162 beats per minute ($0.75 \times 190 = 143$; $0.85 \times 190 = 162$). High-intensity exercise burns an average of 14.3 calories per minute depending on your

weight. Many exercisers will alternate high-intensity workouts into their weekly schedule. These exercises are also helpful to those who have hit plateaus in their exercise routine and have stopped losing weight. If you exercise at a medium-intensity level 4–5 days a week, you may want to try including small amounts of high-intensity exercises. Examples of high-intensity exercises include:

Running

risky climbing stairs

High-level aerobics class

Spinning

Jumping rope

Strength/Weight training(Sharon et al,2011).

Warm Up and Cool Down

Intense exercises should include a warm-up and a cool-down period. Warm-up and cool-down should be 5–10 minutes each, and include a low-level activity such as walking. After warming up and cooling down, it is important to incorporate a series of stretches. Each stretch should be held slowly for 20–30 seconds (without bouncing) and include each major muscle group challenged during the workout (Sharon et al 2011).

Appendix V Session Plan for Twelve Weeks (Three Months) of Experimental Study

1stMonth(October,2020) Training Schedule for Group-II participants.

Duration : 60 min/session

Intensity of exercises: low intensity (~40-50%HRmax) for the first two consecutive weeks, then moderate intensity (~60-65%HRmax) for the rest ten weeks.

Objective: To create physiological adaptation, confidence and awareness.

Day	Types of aerobics training	Time	No Ex	S	Weeks							
					1 st week		2 nd week		3 rd week		4 th week	
					Rep	Rest	Rep	Rest	Rep	Rest	Rep	Rest
Mon	Warming up (walking, jogging etc.)	10m										
	Light stretch	10m	6	1	6x1	30s	6x1	30s	6x2	30s	6x2	30s
	Brisk walking on treadmill speed 5mph level 6%	10m	Wal	1	1	30s	1	30s	1	30s	1	30s
	Cycling	10m	1	1	1	30s	1	30s	1	30s	1	30s
	Sit ups	10m	1	1	1x8	30s	1x8	30s	1x12	30s	1x12	30s
	Cooling down	10m										
Tues	Rest Day											
Wed	Warming up	10m										
	Light stretch	10m	7	1	7x1	30s	7x1	30s	7x2	30s	7x2	30s
	Sit ups	15m	1	1	1x1	30s	1x10	30s	1x15	30s	1x15	30s
	Steps	15m	1	2	1x1	30s	1x15	30s	1x20	30s	1x20	30s
	Cooling down	10m										
Thu	Rest Day											
Fri	Warming up	10m										

	Light stretch	10m	7	1	7x1	30s	7x1	30s	7x1	30s	7x1	30s
	Steps	10m	1	1	1x20	30s	1x20	30s	1x25	30s	5x25	30s
	Running on treadmill	10m	Run	1	1	30s	1	30s	1	30s	1	30s
	Cycling	10m	1	1	1	30s	1	30s	1	30s	1	30s
	Cooling down	10m										
Sat	Rest Day											
Sun	Rest Day											

2nd Month (November,2020) Training Schedule for Group-II participants.

Duration: 60 min/session

Intensity of Exercises: moderate intensity (~65-75%HRmax) for the rest ten weeks.

Objective: For body weight reduction.

Day	Types of aerobics training	Time	No of exe.	Se Ts	Weeks							
					1 st week		2 nd week		3 rd week		4 th week	
					Rep	Rest	Rep	Rest	Rep	Rest	Rep	Rest
Mon	Warming up (walking, jogging etc)	10m										
	Light stretch	10m	8	1	8x1	30s	8x1	30s	8x1	30s	8x1	30s
	Brisk walking on treadmill speed 5mph level 6%	10m	Wal k	1	1	30s	1	30s	1	30s	1	30s
	Cycling	10m	1	2	1	30s	1	30s	1	30s	1	30s

	Sit ups	10m	1	1	1x1 2	30s	1x12	30s	1x1 2	30s	1x1 2	30s
	Cooling down	10m										
Tues	RestDay											
Wed	Warming up	10m										
	Light stretch	10m	6	1	6x3	30s	6x3	30s	6x3	30s	6x3	30
	Sit ups	15m	1	1	1x1 5	30s	1x15	30s	1x1 5	30s	1x1 5	30s
	Steps	15m	1	3	1x1 2	30s	1x12	30s	1x1 2	30s	1x1 2	30s
	Cooling down	10m										
Thu	Rest Day											
Fri	Warming up	10m										
	Light stretch	10m	9	1	9x1	30se c	9x1	30se c	9x1	30s	9x1	30s
	Steps	10m	1	3	1x1 5	30se c	1x15	30se c	1x1 5	30s	1x1 5	30s
	Running on treadmill	10m	Run	1	1	30se c	1	30se c	1	30s	1	30s
	Weight free exercise	10m	4	1	4x6	30se c	4x6	30se c	4x6	30s	4x6	30s
	Cooling down	10m										
Sat	Rest Day											
Sun	Rest Day											

3rdMonth (December,2020) Training Schedule for Group-II participants.

Duration :60 min/session

Intensity of Exercises: moderate intensity (~65-75%HRmax) for the rest ten weeks.

Objective: to body weight reduction.

Day	Types of aerobics training	Time	No of exe.	S e T s	Weeks							
					1 st week		2 nd week		3 rd week		4 th week	
					Rep	Rest	Rep	Rest	Rep	Rest	Rep	Rest
Mon	Warming up (walking, jogging etc.)	10m										
	Weight free exercise	10m	5	3	5x6	30s	5x6	30s	5x6	30s	5x6	30s
	Aerobics dance	20m	Da nce	1	1	30s	1	30s	1	30s	1	30s
	Light stretch	10m	10	2	10x2	30s	10x2	30s	10x2	30s	10x2	30s
	Cooling down	10m										
Tues	RestDay											
Wed	Warming up	10m										
	Brisk walking on treadmill speed 5mph level 6%	20m	1	1	1	30s	1	30s	1	30s	1	30s
	Sit ups	10m	1	1	1x20	30s	1x20	30s	1x20	30s	1x20	30s
	Steps	10m	1	2	1x20	30s	1x20	30s	1x20	30s	1x20	30s
	Cooling	10m										

	down											
Thu	Rest Day											
Fri	Warming up	10m										
	Light stretch	10m	12	2	12x2	30s	12x2	30s	12x2	30s	12x2	30s
	Steps	10m	1	3	15x2	30s	15x2	30s	15x2	30s	15x2	30s
	Aerobics dance	20m	Dance	1	1	30s	1	30s	1	30s	1	30s
	Cooling down	10m										
Sat	Rest Day											
Sun	Rest Day											

Source: Solomon, w. 2010

Appendix V I. Standard Norms of Body Mass Index and Waist Circumference

BMI (kg/m ²)Classification	
< 18.5	Under Weight
18.5 – 24.9	Healthy Weight
25.0 – 29.9	Over Weight
30.0 – 39.9	Obesity

Norm for waist circumference in adults

Waist Circumference		
Risk category	Females	Males
Very low	<27.5 in (< 70cm)	<31.5 in (< 80cm)
Low	27.5-35.0 in(70-89cm)	31.5- 39.0 in (80-99cm)
High	35.5- 43.0 in (90-109cm)	39.5- 47.0 in (100-120cm)
Very high	>43.5 in (> 110 cm)	>47.0 in (>120 cm)

Appendix VII Raw Data Test Recording Sheet

Group I

Control Group										
	B/w		Ht		w/c		Age		BMI	
	Pt	Pot	PT	Pot	Pt	Pot	Pt	Pot	Pt	Pot
S 1	74	75	1.65	1.65	90	91	30	30	27.18	27.54
S 2	71	73	1.63	1.63	92	94	29	29	26.72	27.47
S 3	73	74	1.65	1.65	91	91	28	28	26.81	27.18
S 4	69	70	1.65	1.65	89	90	31	31	25.34	25.71
S 5	71	73	1.67	1.67	93	95	34	34	25.45	26.17
S 6	75	77	1.66	1.66	90	92	30	30	27.21	27.94
S 7	69	69	1.65	1.65	89	90	29	29	25.34	25.34
S 8	70	68	1.61	1.61	89	90	32	32	27.00	26.23
S 9	75	77	1.65	1.65	90	91	30	30	27.54	28.28
S 10	74	75	1.67	1.67	89	90	29	29	26.53	26.89
S 11	71	70	1.65	1.65	89	89	33	33	26.07	25.71
S 12	74	77	1.63	1.63	92	91	27	27	27.85	28.98
S 13	73	73	1.69	1.69	93	93	32	32	25.55	25.55
S 14	72	74	1.66	1.66	94	95	28	28	26.12	26.85
S 15	74	76	1.67	1.67	90	91	34	34	26.53	27.25
S 16	73	75	1.68	1.68	93	94	31	31	25.86	26.57
S 17	72	73	1.65	1.65	89	90	29	29	26.44	26.81
S 18	72	74	1.66	1.66	91	93	30	30	26.12	26.85
S 19	70	70	1.60	1.60	91	92	27	27	27.34	27.34
S 20	71	72	1.61	1.61	92	93	30	30	27.39	27.77

Group II

Experimental Group										
	B/w		Ht		w/c		Age		BMI	
	Pt	Pot	Pt	Pot	Pt	Pot	Pt	Pot	Pt	Pot
S 01	70	66	1.63	1.63	92	85	27	27	26.34	24.84
S 02	73	70	1.64	1.64	90	81	26	26	27.14	26.02
S 03	70	66	1.65	1.65	93	83	30	30	25.71	24.24
S 04	69	65	1.63	1.63	89	81	28	28	25.97	24.46
S 05	74	70	1.68	1.68	93	85	32	32	26.21	24.80
S 06	71	67	1.66	1.66	90	80	30	30	25.76	24.46
S 07	70	65	1.63	1.63	94	83	29	29	26.34	24.46
S 08	72	68	1.68	1.68	91	85	31	31	25.51	24.09
S 09	72	67	1.67	1.67	92	84	29	29	25.81	24.02
S 010	70	66	1.65	1.65	92	86	27	27	25.71	24.24
S 011	69	64	1.60	1.60	93	81	33	33	26.95	25
S 012	70	67	1.62	1.62	89	83	34	34	26.67	25.52
S 013	70	65	1.63	1.63	94	83	34	34	26.34	24.46
S 014	73	69	1.67	1.67	92	82	32	32	26.17	24.74
S 015	70	67	1.60	1.60	89	80	30	30	27.34	26.17
S 016	73	68	1.68	1.68	93	82	34	34	25.86	24.09
S 017	73	68	1.67	1.67	92	90	30	30	26.17	24.38
S 018	69	66	1.60	1.60	91	83	25	25	26.95	25.78
S 019	74	70	1.70	1.70	91	80	35	35	25.60	24.22
S 020	72	69	1.66	1.66	89	81	34	34	26.12	25.03

Pt= pre training, pot= post training, BW = body weight, WC= waist circumference,

HT =height, Ag= age, BMI= Body mass index